

MODULI SPACES OF MEROMORPHIC FUNCTIONS AND DETERMINANT OF THE LAPLACIAN

LUC HILLAIRET, VICTOR KALVIN, AND ALEXEY KOKOTOV

ABSTRACT. The Hurwitz space is the moduli space of pairs (X, f) where X is a compact Riemann surface and f is a meromorphic function on X . We study the Laplace operator $\Delta^{|df|^2}$ of the flat singular Riemannian manifold $(X, |df|^2)$. We define a regularized determinant for $\Delta^{|df|^2}$ and study it as a functional on the Hurwitz space. We prove that this functional is related to a system of PDE which admits explicit integration. This leads to an explicit expression for the determinant of the Laplace operator in terms of the basic objects on the underlying Riemann surface (the prime-form, theta-functions, the canonical meromorphic bidifferential) and the divisor of the meromorphic differential df . The proof has several parts that can be of independent interest. As an important intermediate result we prove a decomposition formula of the type of Burghelaa-Friedlander-Kappeler for the determinant of the Laplace operator on flat surfaces with conical singularities and Euclidean or conical ends. We introduce and study the S -matrix, $S(\lambda)$, of a surface with conical singularities as a function of the spectral parameter λ and relate its behavior at $\lambda = 0$ with the Schiffer projective connection on the Riemann surface X . We also prove variational formulas for eigenvalues of the Laplace operator of a compact surface with conical singularities when the latter move.

1. INTRODUCTION

1.1. General part. Studying the determinants of Laplacians on Riemann surfaces is motivated by needs of quantum field theory (in connection with various partition functions) and geometric analysis (in particular, in connection with the Sarnak program, [36]). The explicit expressions for the determinant of the Laplacian in the metric of constant negative curvature ([7]) and in the Arakelov metric (obtained in [2] in relation to so-called bosonization formulas from string theory) for compact Riemann surfaces of genus $g > 1$ are among the most spectacular results of the subject. According to the Sarnak program, these determinants (which are functions on the moduli space of Riemann surfaces) can be used to study the geometry of the moduli space via methods of Morse theory. In particular, their behavior at the boundary of moduli space is of great importance and was intensively studied (see, e.g., [42], [41]).

It seems very interesting to consider the case which is in a certain sense opposite to the case of the metric of constant curvature: instead of distributing the curvature uniformly along the Riemann surface X one can concentrate it at a finite set $\{P_1, \dots, P_M\} \subset X$. This leads to a flat metric \mathbf{m} on X with singularities (e.g.,

Received by the editors August 11, 2016.

2010 *Mathematics Subject Classification.* Primary 58J52, 47A10, 30F30; Secondary 14H15, 14H81, 14K25.

conical) at P_k . Determinants of Laplacians for various classes of singular flat metrics were introduced and studied at least on the formal level (via path integrals) by physicists ([38], [43], [20], [3]) and certain explicit expressions for them were produced (see, e.g., [38], [43]).

One of the main challenges is to study such determinants from the spectral theory of self-adjoint operators and perturbation theory points of view. This was done in the mathematical literature for the determinants of the Laplacians of smooth metrics; in particular, those two mentioned above (see, e.g., Fay's book [9] for complete compendium and consistent exposition). The standard definition of the determinant uses the ζ -function of the corresponding Laplace operator

$$(1.1) \quad \ln \det \Delta^{\mathbf{m}} = -\zeta'_{\Delta^{\mathbf{m}}}(0),$$

$$(1.2) \quad \zeta_{\Delta^{\mathbf{m}}}(s) = \sum_j \frac{1}{\lambda_j^s},$$

where in the latter expression the sum is extended over all non-zero eigenvalues of $\Delta^{\mathbf{m}}$. This definition makes sense when the metric \mathbf{m} has conical singularities provided a regular self-adjoint extension is considered; for instance, the Friedrichs one (see [15] for the definition of regular). Indeed, the Laplace operator $\Delta^{\mathbf{m}}$ (with natural domain consisting of smooth functions on X supported outside the conical points P_k) is not essentially self-adjoint (cf. [19]). Comparing the determinants of the different self-adjoint extensions of $\Delta^{\mathbf{m}}$ leads to a nice application of Birman-Krein theory and is done in [15] (see also the references therein). In what follows we consider the Friedrichs extension of $\Delta^{\mathbf{m}}$; our results refer to this self-adjoint extension only.

In [25] an explicit expression was found for the determinant of (the Friedrichs extension of) the Laplace operator corresponding to a flat conical metric \mathbf{m} with *trivial holonomy*. Any metric of this type can be written $|\omega|^2$, where ω is a holomorphic one-form on X , zeros of ω of multiplicity ℓ are the conical points of the metric $|\omega|^2$ with conical angle $2\pi(\ell+1)$. The moduli space of pairs (X, ω) , where X is a compact Riemann surface and ω is a holomorphic one-form on X , is stratified according to the multiplicities of ω (see [29]). In [25] it was proved that on each stratum of the moduli space of holomorphic differentials the ratio

$$\frac{\det \Delta^{|\omega|^2}}{\text{Area}(X) \det \Im \mathbb{B}},$$

where \mathbb{B} is the matrix of b -periods of the Riemann surface X , coincides with the modulus square of a holomorphic function τ on the stratum. This holomorphic function τ (the so-called Bergman tau-function on the space of holomorphic differentials) admits explicit expression through theta-functions, prime-forms, and the divisor of the holomorphic one-form ω . In the case $g = 1$ the holomorphic one-forms have no zeros, the metric $|\omega|^2$ is smooth, and the corresponding result coincides with the classical Ray-Singer formula for the determinant of the Laplacian on an elliptic curve with flat conformal metrics.

In [23] a comparison formula (an analog of the classical Polyakov formula) relating determinants of the Laplacians in two conformally equivalent flat conical metrics was found. This led to the generalization of the results of [25] to the case of arbitrary flat conformal metrics with conical singularities.

Together with determinant of the Laplacians in flat conical metrics given by the modulus square of the holomorphic one-form (these metrics have finite volume and the spectra of the corresponding self-adjoint Laplacians are discrete) in physics literature appear determinants of the Laplacians corresponding to flat metrics $|\omega|^2$, where ω is now a *meromorphic* one-form on X . Depending on the order of the poles of ω , the corresponding non-compact Riemannian manifold $(X, |\omega|^2)$ of infinite volume has cylindrical, Euclidean, or conical ends. The spectrum of the corresponding Laplace operator is continuous (with possible embedded eigenvalues, say, in case of cylindrical ends) and the Ray-Singer regularization of the determinant ((1.1), (1.2)) is no longer applicable. The way to regularize such determinants is, in principle, also well known (see, e.g., [32]): considering the Laplacian Δ as a perturbation of some properly chosen “free” operator $\mathring{\Delta}$, one introduces a relative determinant $\det(\Delta, \mathring{\Delta})$ in terms of the relative ζ -function

$$(1.3) \quad \zeta(s; \Delta, \mathring{\Delta}) = \frac{1}{\Gamma(s)} \int_0^\infty \text{Tr}(e^{-\Delta t} - e^{-\mathring{\Delta} t}) t^{s-1} dt,$$

where a suitable regularization of the integral is made (being understood in the conventional sense the integral is usually divergent for any value of s).

Following this approach, in [16] we studied the regularized determinants

$$\det(\Delta, \mathring{\Delta}) = e^{-\zeta'(0; \Delta, \mathring{\Delta})}$$

of the Laplacians on the so-called Mandelstam diagrams - the flat surfaces with cylindrical ends (more precisely, Riemann surfaces X with the metric $|\omega|^2$, where ω is a meromorphic one-form on X having only simple poles and such that all the periods of ω are pure imaginary and all the residues of ω at the poles are real).

In the present paper we consider determinants of the Laplacian corresponding to flat metrics with even wilder singularities: the corresponding Riemannian manifolds have Euclidean (i.e., isometric to a vicinity of the point at infinity of the Euclidean plane) and/or conical ends (i.e., isometric to a vicinity of the point at infinity of a straight cone).¹

¹As is well known in scattering theory, the spectral properties change drastically depending on the structure of the ends, and, in particular, the techniques that can be employed to define a spectral determinant and then to prove the BFK gluing formula. More precisely, for low energies, in the cylindrical case the (absolutely) continuous spectrum of the Laplacian is of *finite multiplicity* and thus everything can be understood by eventually considering a finite dimensional perturbation (roughly speaking one only has to consider what happens on the $0 - th$ mode for each cylindrical end), whereas in the case of Euclidean or conical ends the continuous spectrum is of *infinite multiplicity* which raises the problem to a very different level. Much more involved analysis has to be performed first, in order to define the determinant, and then to prove the BFK formula.

Thus by using a conventional approach to the definition of a relative determinant and to the proof of the BFK formula, one intimately deals with analysis of the essential singularity of the resolvent at energy zero (thanks to continuous spectrum of infinite multiplicity), which seems to be a very challenging problem. In particular, it is highlighted by the fact that due to exactly the same difficulty the assumption $\ker D_Y = \{0\}$ in [33] is not yet removed after years of attempts.

We should also note that in [14] the authors define the determinant and prove a BFK formula for the Laplacian with a similar spectral portrait. Unfortunately, despite their approach of avoiding studying the resolvent essential singularity, it fails in our geometric settings.

In the present paper we use a different approach. Developing further some ideas due to Carron, we base our analysis on the study of the Neumann jump operator instead of studying essential singularity of the resolvent at energy zero. We then obtain all needed information on spectral shift functions thanks to a relation between zeta-determinant of the Neumann jump operators

These metrics are given as the modulus square of the differential of an arbitrary meromorphic function f on a compact Riemann surface X . The moduli space of pairs (X, f) is called the Hurwitz space \mathcal{H} . We define and study the regularized determinant of the Laplace operator corresponding to the metric $|df|^2$ as a functional on \mathcal{H} . The main result of this work is an explicit formula for the determinant.

It should be mentioned that such determinants appeared for the first time in [43], [3] (see also [20]), although no attempt was made to define them rigorously.

1.2. Results and organization of the paper. Let X be a Riemann surface and let f be a meromorphic function $f : X \rightarrow \mathbb{P}^1$. The metric $|df|^2$ gives to X a structure of a (non-compact) flat Riemannian manifold with conical singularities and conical (or Euclidean) ends. The conical singularities are located at the critical points P_1, \dots, P_M of f (or, equivalently, at the zeros of the meromorphic one-form df), the ends of X are located at the poles of df . The moduli space of (equivalence classes) of such pairs (X, f) is known as Hurwitz space \mathcal{H} and the critical values $z_m, z_m = f(P_m)$, $m = 1, \dots, M$, locally parametrize \mathcal{H} .

Given such a Riemannian manifold $(X, |df|^2)$, we introduce the reference manifold $(\hat{X}, \hat{\mathbf{m}})$ as the disjoint union of the complete cones corresponding to the ends of $(X, |df|^2)$. By Δ and $\hat{\Delta}$ we denote the Laplace operators on $(X, |df|^2)$ and $(\hat{X}, \hat{\mathbf{m}})$ correspondingly.

The first part of the paper aims at defining the relative zeta-regularized determinant $\det_\zeta(\Delta, \hat{\Delta})$ and proving a version of the Burghelca-Friedlander-Kappeler (BFK in what follows) gluing formula (see [4]). This new BFK type formula is a generalization of the Hassell-Zelditch formula for the determinant of the Laplacian in exterior domains [14]; here we rely on ideas from [5, 6, 14].

In order to obtain the gluing formula, we cut X along some hypersurface Σ . This decomposes X into a compact part X_- and the union of conical/Euclidean ends X_+ . The latter is isometric to the reference surface \hat{X} with a compact part \hat{X}_- removed. There is some latitude in choosing the initial Σ . In order to choose Σ we first specify some large R and then in each end of $(X, |df|^2)$ we take a circle whose radius depends on R and on the cone angle of the end; see Definition 3. As expected the gluing formula then involves the Neumann jump operator \mathcal{N} on Σ and the Dirichlet Laplacian Δ_-^D on X_- ; see Theorem 1 below.

Theorem 1. *For R large enough we have the BFK gluing formula*

$$\det_\zeta(\Delta, \hat{\Delta}) = C \det_\zeta^* \mathcal{N} \cdot \det_\zeta \Delta_-^D,$$

where \mathcal{N} , Δ_-^D depend on R . The constant C depends on R but not on the moduli parameters z_1, \dots, z_M as long as the corresponding critical points P_m do not approach Σ .

Note that the proof of the gluing formula also holds for a more general class of metric (see Remark 2).

and spectral shift functions. Let us also note that in [16] we rely on the conventional approach of studying the resolvent at zero energy.

Let us now sketch some steps leading to this theorem. First we start from the BFK gluing formula for $\det_{\zeta}(\Delta - \lambda, \mathring{\Delta} - \lambda)$ obtained in [5] for negative (regular) values of the spectral parameter λ . In order to obtain a gluing formula for $\det_{\zeta}(\Delta, \mathring{\Delta})$, we study the behavior of all ingredients in the gluing formula for $\det_{\zeta}(\Delta - \lambda, \mathring{\Delta} - \lambda)$ as $\lambda \rightarrow 0-$ (i.e., at the bottom of the continuous spectrum of Δ and $\mathring{\Delta}$) and then pass to the limit. As usual, this essentially reduces to derivation of asymptotics as $\lambda \rightarrow 0-$ for the zeta-regularized determinant of the Neumann jump operator and for the spectral shift function of the pair $(\Delta, \mathring{\Delta})$. In principle, both asymptotics were obtained in [6] for Schrödinger type operators on manifolds with conical ends. Unfortunately, those asymptotics cannot be used for our purposes because the asymptotic for the Neumann jump operator contains an unspecified constant and the asymptotic for the spectral shift function is not sufficiently sharp. We demonstrate that at least in our setting (no potential) the methods of [6] can be improved to specify the constant and to obtain a sufficiently sharp asymptotic of the spectral shift function as needed for the proof of our BFK formula. Once these asymptotics are obtained, we follow the lines of [14] in our study of the behavior of $\det_{\zeta}(\Delta - \lambda, \mathring{\Delta} - \lambda)$ as $\lambda \rightarrow 0-$ and also in definition of $\det_{\zeta}(\Delta, \mathring{\Delta})$.

Using this BFK formula, we prove (as it was done in similar situations in [23], [16]) that the variations of the determinant of the Laplacian with respect to the moduli parameters z_k remain the same if we replace the metric $\mathbf{m} = |df|^2$ of infinite volume by a metric $\tilde{\mathbf{m}}$ of finite volume, where $\tilde{\mathbf{m}}$ coincides with \mathbf{m} outside vicinities of the poles of f and with some standard non-singular metric of finite volume inside these vicinities. The aim of the second part of the paper is thus to study the zeta-regularized determinant of this new metric $\tilde{\mathbf{m}}$ and its variation with respect to moduli parameters.

We show that these variations can be conveniently expressed using the so-called S -matrix, so we start the second part of the paper by introducing this object and deriving some of its properties. We think that the S -matrix is an important characteristic of a compact Riemann surface X equipped with a conformal metric $\tilde{\mathbf{m}}$ with conical singularities. It is introduced in analogy with scattering problems and the general theory of boundary triples (see [13]). Basically, the S -matrix is a meromorphic matrix function of the spectral parameter λ . It serves as a formal analog of the scattering matrix on complete non-compact manifolds (say, on manifolds with cylindrical ends) in the sense that the elements of S are coefficients in asymptotics of certain eigenfunctions (all growing terms in the asymptotics near conical singularities of $\tilde{\mathbf{m}}$ are interpreted as incoming waves and all decaying terms — as outgoing). Let us also note that there is no true scattering on the (incomplete) manifold $(X, \tilde{\mathbf{m}})$ (the corresponding operators have no continuous spectrum) and that $S(\lambda)$ is a non-unitary matrix with poles at the eigenvalues λ of the Friedrichs self-adjoint Laplacian on $(X, \tilde{\mathbf{m}})$ [15].

It turns out that the value, $S(0)$, of the S -matrix can be found explicitly: it depends only on the conical angle at the conical point, the conformal class of the surface X , and the choice of the holomorphic local parameter near the conical point (the so-called distinguished local parameter for the metric $\tilde{\mathbf{m}}$). For instance, when the conical angle is 4π , we express the matrix elements of $S(0)$ through the Bergman reproducing kernel for holomorphic differentials and a certain special projective connection on X (the so-called Schiffer projective connection).

In the general case we prove that a certain linear combination of the matrix elements of $S(0)$ (actually, the one that appears later in the variational formulas for the determinant) can be expressed as the derivative (of the order depending on the conical angle) of the Schiffer projective connection.

We continue by studying the moduli variations of the zeta-regularized determinant of $\Delta^{\tilde{\mathbf{m}}}$. We use the Kato-Rellich perturbation theory to compute the variation of individual eigenvalue branches and then a contour argument similar to the one from [15] to get the variational formula for the determinant. This formula involves a combination of the matrix elements of $S(0)$ and hence the Schiffer projective connection. Writing it into an invariant form, we obtain the following theorem.

Theorem 2. *Let P_m be a zero of the meromorphic differential df of multiplicity ℓ_m and let $z_m = f(P_m)$ be the corresponding critical value of f . Let the metric $\tilde{\mathbf{m}}$ be obtained from the metric $|df|^2$ via smoothing of the conical ends. Then*

$$(1.4) \quad \partial_{z_m} \ln \frac{\det_{\zeta}^*(\Delta^{\tilde{\mathbf{m}}})}{\det \Im \mathbb{B}} = -\frac{1}{12\pi i} \oint_{P_m} \frac{S_B - S_f}{df},$$

where S_B is the Bergman projective connection, $S_f = \frac{f'''f' - \frac{3}{2}(f'')^2}{(f')^2}$ is the Schwarzian derivative, and \mathbb{B} is the matrix of b -periods.

In this theorem we can replace $\det_{\zeta}^*(\Delta^{\tilde{\mathbf{m}}})$ by $\det(\Delta, \mathring{\Delta})$ since we proved before that the moduli variations of both functions coincide.

The system of PDE for $\det \Delta^{\tilde{\mathbf{m}}}$ that appears in Theorem 2 is the governing system for the Bergman tau-function on the Hurwitz space (introduced and studied in [22], [25], [27], [26]). The latter system was explicitly integrated in [24] and in §5 we remember this result (unfortunately, technically involved). This leads to the following explicit formula for $\det_{\zeta}(\Delta, \mathring{\Delta})$.

Theorem 3. *Let (X, f) be an element of the Hurwitz space $\mathcal{H}(M, N)$ and let $\tau(X, f)$ be given by expressions ((6.86), (6.85), (6.84)). There is the following explicit expression for the regularized relative determinant of the Laplacian Δ on the Riemann surface X :*

$$(1.5) \quad \det_{\zeta}(\Delta, \mathring{\Delta}) = C \det \Im \mathbb{B} |\tau|^2,$$

where C is a constant that depends only on the connected component of the space $\mathcal{H}(M, N)$ containing the element (X, f) .

We finish the paper with two illustrating examples in genus 0, deriving the formulas for the determinant of the Laplacian on the space of polynomials of degree N and on the space of rational functions with three simple poles.

2. THE REGULARIZED DETERMINANT AS A FUNCTIONAL ON THE HURWITZ SPACE AND A BFK GLUING FORMULA

To a pair (X, f) , where X is a compact Riemann surface and f is a meromorphic function on X (i.e., to an element of the Hurwitz space), there corresponds a Riemannian manifold $(X, |df|^2)$. Our aim is to define a regularized determinant of the corresponding Laplacian and to prove a BFK type gluing formula. Since the metric $\mathbf{m} = |df|^2$ has conical singularities and non-compact conical ends, this is not that straightforward and requires several steps. First, we consider regular values of the spectral parameter λ^2 , i.e., $\lambda^2 \in \mathbb{C} \setminus [0, \infty)$. In that case, the definition of

the relative determinant and the BFK gluing formula are the same as in [5]. Then we derive estimates for the determinant of the Dirichlet-to-Neumann operator as λ approaches 0; our methods here are closely related to those of [6]. These estimates allow us to define a zeta-regularized determinant for $\lambda = 0$ similarly to [14]. Then we prove the gluing formula for the thus defined determinant. At the end of this section we use the gluing formula to compactify $(X, |df|^2)$ in such a way, that locally, the moduli variations remain the same.

2.1. The flat Laplacian of an element in the Hurwitz space. We will be dealing with conical singularities and conical ends. These are defined in the following way.

Definition 1.

- For any $\ell \in \mathbb{N}$ the Euclidean cone of total angle $2\ell\pi$ is the Riemannian manifold $(\mathbb{C}, |\ell y^{\ell-1} dy|^2)$.
- A point P in a Riemannian manifold will be a conical singularity of angle $2\ell\pi$ if there is a neighborhood of P that is isometric to the set $(\{|y| < \varepsilon\}, |\ell y^{\ell-1} dy|^2)$ for some positive ε .
- An open set $\Omega \subset X$ of a Riemannian manifold (X, \mathbf{m}) such that (Ω, \mathbf{m}) is isometric to $(\{|y| > R\}, |\ell y^{\ell-1} dy|^2)$ for some positive R will be called a conical end of angle $2\ell\pi$ (Euclidean end if $\ell = 1$).

Let (X, \mathbf{m}) be a Riemannian manifold such that the metric is flat with a finite number of conical singularities and conical ends. Denote by Δ the self-adjoint Friedrichs extension of the (non-negative) symmetric Laplace operator defined on smooth compactly supported functions that vanish near the conical singularities.

Let f be a meromorphic function on a compact Riemann surface X of genus $g \geq 0$ or, what is the same, a ramified covering of the Riemann sphere

$$(2.6) \quad f : X \rightarrow \mathbb{P}^1.$$

Two coverings $f_1 : X_1 \rightarrow \mathbb{P}^1$ and $f_2 : X_2 \rightarrow \mathbb{P}^1$ are called equivalent if there exists a biholomorphic map $g : X_1 \rightarrow X_2$ such that $f_1 = f_2 \circ g$.

The following constructions are standard; we recall them for the convenience of the reader.

The critical points, $P_m, m = 1, \dots, M$, of the function f (i.e., those points for which $df(P_m) = 0$) are the ramification points of the covering, the points $z_m = f(P_m)$ are called the critical values. The ramification index of the covering at the point P_m equals $\ell_m + 1$, where ℓ_m is the order of the zero of the one-form df at P_m . Denote by $\infty_1, \dots, \infty_K$ the poles of f , and let k_1, \dots, k_K be their multiplicities.

Then the covering (2.6) has degree $N = k_1 + \dots + k_K$ and the following Riemann-Hurwitz formula holds:

$$\sum_{m=1}^M \ell_m - \sum_{j=1}^K (k_j + 1) = 2g - 2,$$

where g is the genus of X .

Pick some regular value $z_0 \in \mathbb{P}^1$ and draw on \mathbb{P}^1 the segments $I_0 := [z_0, \infty]$, $I_m = [z_0, z_m], m = 1, \dots, M$. It may happen that some segment is repeated several times if different critical points take the same critical value. We may also choose z_0 such that all these segments have pairwise disjoint interiors. Denote by $L := \bigcup_{m=0}^M I_m$ the union of these segments and observe that $\mathbb{P}^1 \setminus L$ contains only regular values of

f . It follows that $X \setminus f^{-1}(L)$, the complement of the preimage of L by f in X has N connected components. By construction f is a biholomorphic map from each of these connected components onto $\mathbb{P}^1 \setminus L$. We denote these connected components by $\mathbb{C}_n, n = 1 \dots, N$ and call them the sheets of the covering. Each \mathbb{C}_n can be seen as a copy of the complex plane equipped with the cuts provided by L .

On each sheet, the metric $|dz|^2$ lifted from the base \mathbb{P}^1 to the covering space coincides with the metric $|df|^2$. The Riemannian manifold $(X, |df|^2)$ is thus obtained by gluing N copies of a Euclidean plane $(\mathbb{C}, |dz|^2)$ with a system of non-intersecting cuts, one of which extends to infinity.

For each critical point P_m of ramification index $\ell_m + 1$, we obtain an $(\ell_m + 1)$ -cycle γ_j obtained by looking to see in which order the sheets are following one another when making a small loop around P_m . It follows that P_m is a conical singularity of angle $2\pi(\ell_m + 1)$.

For each $z_m, m \neq 0$, we obtain a permutation in S_N by composing the cycles for each critical point in $f^{-1}(z_m)$. We thus obtain M' permutations $\sigma_{m'}, m' = 1 \dots M'$, where M' is the number of different critical values. The critical value z_m is thus associated with one cycle in one of the permutations $\sigma_{m'}, m' = 1 \dots M'$.

In the same manner we obtain a permutation σ_0 by looking at the preimage of a large loop that surrounds z_0 (or, equivalently, a small loop around ∞ in the base \mathbb{P}^1). This permutation describes the structure at infinity of the Riemannian manifold $(X, |df|^2)$: each fixed point of σ_0 corresponds to a flat Euclidean end and a cycle of length k to a conical end of angle $2k\pi$. A pole in f of order k corresponds to a conical end of angle $2k\pi$ (and, therefore, a Euclidean end for a simple pole).

The flat structure on $(X, |df|^2)$ is completely characterized by the positions of the critical values $z_m, m = 1 \dots M$, and by the permutations $\sigma_{m'}, m' = 0 \dots M'$. Conversely, starting from $M' + 1$ permutations of S_N , and M' distinct points $w_1, \dots, w_{m'}$ in \mathbb{C} , we construct the sequence z_0, \dots, z_M by choosing a distinct point z_0 and, for $m > 0$, by repeating $w_{m'}$ as many times as there are disjoint cycles in $\sigma_{m'}$. We then glue N sheets according to the scheme prescribed by the permutations and obtain a (not necessarily connected) flat surface (X, \mathbf{m}) with conical singularities and conical ends.

It turns out that it is always possible to find a meromorphic function f from X to \mathbb{P}^1 such that (X, \mathbf{m}) is isometric to $(X, |df|^2)$.

Introduce the Hurwitz space $\mathcal{H}(N, M)$ of equivalence classes of coverings $f : X \rightarrow P^1$ of degree N with M ramification points of (fixed) indices $\ell_1 + 1, \dots, \ell_M + 1$ and K poles of (fixed) multiplicities $k_1, \dots, k_K; k_1 + \dots, k_K = N$. The space $\mathcal{H}(N, M)$ is a complex manifold of dimension M (see [11]; we notice here that it may have more than one connected component) and the critical values z_1, \dots, z_M can be taken as local coordinates on $\mathcal{H}(N, M)$.

If all the critical points of the maps f are simple, then the corresponding Hurwitz space is usually denoted by $\mathcal{H}_{g,N}(k_1, \dots, k_K)$ and is known to be connected (see [34]).

Definition 2. We will refer to the coordinates z_1, \dots, z_M as *moduli*.

From the flat metric point of view, moving z_m can be easily realized by cutting a small ball around P_m ; then move P_m inside this ball. Since the boundary of the ball does not change we can glue the new ball back into the surface.

For such a Riemannian manifold $(X, |df|^2)$ we define a reference manifold $(\mathring{X}, \mathring{\mathbf{m}})$ which is obtained in the following way. Take those N sheets with cuts that correspond to X , and in the gluing scheme of X , keep σ_0 and replace all the permutations $\sigma_{m'}, m' > 0$ by the identity. It can be easily seen that $(\mathring{X}, \mathring{\mathbf{m}})$ is the disjoint union of cones; those cones correspond to the conical ends of X and the tip of each cone is now located above z_0 .

The Laplacian Δ can be considered as a perturbation of the free Laplacian $\mathring{\Delta} := \Delta^{\mathring{\mathbf{m}}}$ acting in $L^2(\mathring{X})$. The perturbation is basically reduced to the change of the domain of the unbounded operator: when we make slits on \mathring{X} and glue them according to a certain gluing scheme, it induces boundary conditions on the sides of the cuts. The determinant of Δ will then be defined in terms of the relative zeta-function (1.3) as a regularized relative determinant $\det_{\zeta}(\Delta, \mathring{\Delta})$.

The main goal of this work is to study the relative determinant $\det_{\zeta}(\Delta, \mathring{\Delta})$ as a functional on the space $\mathcal{H}(N, M)$.

2.2. Relative determinant and BFK gluing formula for negative energies.

Let X be a compact Riemann surface and let f be a non-constant meromorphic function on X . Introduce the flat singular metric $\mathbf{m} = |df|^2$ on X . As it is explained in the previous section, the flat singular Riemannian manifold (X, \mathbf{m}) has conical points (at the zeros, P_1, \dots, P_M , of the differential df) and conical ends of angle $2\pi k_j$ at the poles, $\infty_1, \dots, \infty_K$, of f , where k_j is the order of the corresponding pole. Let Δ be the (Friedrichs) Laplacian on (X, \mathbf{m}) .

Let $(\mathring{X}, \mathring{\mathbf{m}})$ be the reference unperturbed manifold and let $\mathring{\Delta}$ be the associated Laplace operator. We recall that $(\mathring{X}, \mathring{\mathbf{m}}) = \bigcup_{j=1}^K (\mathbb{C}, |k_j y_j^{k_j-1} dy_j|^2)$.

Since (X, \mathbf{m}) and $(\mathring{X}, \mathring{\mathbf{m}})$ are isometric outside a compact region the methods and results of [6] apply.

For $R > 0$ large enough, there is a subset $X_+(R) \subset X$ that is isometric to

$$(2.7) \quad \bigcup_{j=1}^K \{y_j \in \mathbb{C} : |y_j| \geq R^{1/k_j}\} \subset \mathring{X}.$$

Definition 3. We denote by Σ_R the boundary of the region $X_+(R)$. It is the union of K circles $\{y \in \mathbb{C} : |y| = R^{1/k_j}\}$ on X .

Note that R will be chosen at the very beginning of construction and will then be fixed. In what follows we omit the reference to R and simply write Σ, X_+ .

We represent X in the form

$$X = X_- \cup_{\Sigma} X_+,$$

where $X_- = X \setminus (X_+ \cup \Sigma)$.

Following [5] we first define the external Dirichlet-to-Neumann operator. We consider each conical end $\{y_j \in \mathbb{C}_j : |y_j| \geq R^{1/k_j}\}$ of X_+ separately and omit the subscript j for brevity of notation. We introduce the coordinates (r, φ) where $r = |y|^k \in [R, \infty)$ and $\varphi = \arg y \in (-\pi, \pi]$. We have

$$g = dr^2 + k^2 r^2 d\varphi^2, \quad \Delta = r^{-2}((r\partial_r)^2 + k^{-2}\partial_{\varphi}^2).$$

Separation of variables shows that for $\lambda \in \mathbb{C} \setminus \{0\}$ with $\Im \lambda \geq 0$ the exterior Dirichlet problem

$$(2.8) \quad (\Delta - \lambda^2)u(\lambda) = 0 \text{ on } X_+, \quad u(\lambda) = f \text{ on } \Sigma,$$

has a unique solution of the form

$$u(r, \varphi; \lambda) = \sum_{n=-\infty}^{\infty} C_n \frac{H_{\nu_n}^{(1)}(\lambda r)}{H_{\nu_n}^{(1)}(\lambda R)} e^{in\varphi}, \quad \nu_n = \frac{|n|}{kR},$$

where $f \in C^\infty(\Sigma)$, $C_n = (2\pi)^{-1} \int_{-\pi}^{\pi} f(\varphi) e^{-in\varphi} d\varphi$, and $H_n^{(1)}$ is the Hankel function. This solution is in $L^2(X_+)$ if $\Im\lambda > 0$. If $\Im\lambda = 0$, it is the unique outgoing solution that satisfies the Sommerfeld radiation condition

$$\sqrt{r}(\partial_r u(\lambda) - i\lambda r u(\lambda)) \rightarrow 0 \text{ as } r \rightarrow \infty.$$

The external Dirichlet-to-Neumann operator on Σ acts by the formula

$$(2.9) \quad \mathcal{N}_+(\lambda) f = -\partial_r u(\lambda) \upharpoonright_{r=R}.$$

Thus $\psi_n(\varphi) = (\text{Vol } \Sigma)^{-1/2} e^{in\varphi}$ are eigenfunctions of $\mathcal{N}_+(\lambda)$ with $\|\psi_n\|_{L^2(\Sigma)} = 1$, and

$$(2.10) \quad \mu_n(\lambda) = \mu_{-n}(\lambda) = -\frac{\partial_r H_{\nu_n}^{(1)}(\lambda r) \upharpoonright_{r=R}}{H_{\nu_n}^{(1)}(\lambda R)}$$

are the corresponding eigenvalues (if $\Im\lambda \geq 0$, $\lambda \neq 0$).

We can also consider Σ and X^+ as subsets of \mathring{X} . Then in the same manner we have $\mathring{X} = \mathring{X}_- \cup_{\Sigma} \mathring{X}_+$.

Let Δ_{\pm}^D be the Friedrichs extensions of the Dirichlet Laplacians in $L^2(X_{\pm})$. We denote by $\Delta^D := \Delta_-^D \oplus \Delta_+^D$ the Friedrichs Laplace operator on $L^2(X)$ with Dirichlet boundary condition on Σ . Similarly, we define $\mathring{\Delta}_{\pm}^D$ and $\mathring{\Delta}^D$.

The spectrum $\text{spec}(\Delta^D)$ of the positive self-adjoint operator Δ^D is discrete. For any $\lambda^2 \in \mathbb{C} \setminus \sigma(\Delta^D)$ and $f \in H^1(\Sigma)$ there exists a unique solution $u(\lambda) \in H^{3/2}(X_-)$ to the Dirichlet problem

$$(2.11) \quad (\Delta - \lambda^2)u(\lambda) = 0 \text{ on } X_- \setminus \Sigma, \quad u(\lambda) = f \text{ on } \Sigma,$$

such that

$$(2.12) \quad u(\lambda) = \tilde{f} - (\Delta_-^D - \lambda^2)^{-1}(\Delta - \lambda^2)\tilde{f},$$

where $\tilde{f} \in H^{3/2}(X_-)$ is a continuation of f and

$$(\Delta_-^D - \lambda^2)^{-1} : H^{-1/2}(X_-) \rightarrow H^{3/2}(X_-)$$

is a holomorphic function of $\lambda^2 \in \mathbb{C} \setminus \sigma(\Delta^D)$; here

$$\|u; H^s(X_-)\| = \|(\Delta_-^D)^{s/2} u; L^2(X_-)\|.$$

The Dirichlet-to-Neumann operator $\mathcal{N}_-(\lambda)$ on Σ acts by the formula

$$\mathcal{N}_-(\lambda) f = \partial_r u(\lambda) \upharpoonright_{r=R},$$

where f is the right hand side in (2.11) and $u(\lambda)$ is defined by (2.12). The function $\lambda^2 \mapsto \mathcal{N}_-(\lambda) \in \mathcal{B}(H^1(\Sigma), L^2(\Sigma))$ is holomorphic in $\mathbb{C} \setminus \sigma(\Delta^D)$; here and elsewhere $\mathcal{B}(X, Y)$ stands for the space of bounded operators from X to Y .

Finally, we introduce the Neumann jump operator

$$\mathcal{N}(\lambda) = \mathcal{N}_+(\lambda) + \mathcal{N}_-(\lambda),$$

which is a first order elliptic classical pseudodifferential operator on Σ with the principal symbol $2|\xi|$.

For $\lambda^2 \leq 0$ the operator $\mathcal{N}(\lambda)$ is formally self-adjoint and non-negative, it is positive if $\lambda^2 < 0$, and $\ker \mathcal{N}(0) = \{c \in \mathbb{C}\}$ (see, e.g., [6, Sec. 3.3] for details). (Note that in Theorem 4 the operator $\mathcal{N}(0)$ is denoted by \mathcal{N} .) Let $\lambda^2 < 0$. The function $\zeta(s) = \text{Tr } \mathcal{N}(\lambda)^{-s}$ is holomorphic in $\{s \in \mathbb{C} : \Re s > 1\}$ and admits a meromorphic continuation to \mathbb{C} with no pole at zero; we set $\det_{\zeta} \mathcal{N}(\lambda) = e^{-\zeta'(0)}$.

It is known (see [5, Theorem 2.2]) that the difference

$$(\Delta + 1)^{-1} - (\Delta^D + 1)^{-1}$$

is in the trace class. By the Krein theorem, see, e.g., [40, Chapter 8.9] or [5, Theorem 3.3], there exists a spectral shift function $\xi(\cdot; \Delta, \Delta^D) \in L^1(\mathbb{R}_+, (1+\lambda^2)^{-2} \lambda d\lambda)$ such that

$$(2.13) \quad \text{Tr}((\Delta + 1)^{-1} - (\Delta^D + 1)^{-1}) = - \int_0^\infty \xi(\lambda; \Delta, \Delta^D) (1 + \lambda^2)^{-2} 2\lambda d\lambda.$$

Moreover, the following representation is valid:

$$(2.14) \quad \text{Tr}(e^{-t\Delta} - e^{-t\Delta^D}) = -t \int_0^\infty e^{-t\lambda^2} \xi(\lambda; \Delta, \Delta^D) 2\lambda d\lambda,$$

which implies that the left hand side in (2.14) is absolutely bounded uniformly in $t > \epsilon > 0$. The heat trace asymptotic

$$(2.15) \quad \text{Tr}(e^{-t\Delta} - e^{-t\Delta^D}) \sim \sum_{j \geq -2} a_j t^{j/2}, \quad t \rightarrow 0+,$$

can be obtained in a usual way; see, e.g., [16, Lemma 4]. Thus for $\lambda^2 < 0$ the relative zeta-function given by

$$\zeta(s; \Delta - \lambda^2, \Delta^D - \lambda^2) = \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} e^{\lambda^2 t} \text{Tr}(e^{-t\Delta} - e^{-t\Delta^D}) dt$$

is defined for $\Re s > 1$ and continues meromorphically to the complex plane with no pole at $s = 0$ by the usual argument. The relative determinant is defined to be

$$\det_{\zeta}(\Delta - \lambda^2, \Delta^D - \lambda^2) = e^{-\zeta'(0; \Delta - \lambda^2, \Delta^D - \lambda^2)}.$$

By [5, Theorem 4.2] we have the gluing formula

$$(2.16) \quad \det_{\zeta}(\Delta - \lambda^2, \Delta^D - \lambda^2) = \det_{\zeta} \mathcal{N}(\lambda), \quad \lambda^2 < 0.$$

(Although only smooth manifolds are considered in [5], it is fairly straightforward to see that the argument in [5] remains valid for (2.16) as far as we consider only Friedrichs extensions and there are no conical points on Σ .)

All the constructions above can also be done for $(\mathring{X}, \mathring{\mathbf{m}})$. Thus similarly to (2.16) we have

$$(2.17) \quad \det_{\zeta}(\mathring{\Delta} - \lambda^2, \mathring{\Delta}^D - \lambda^2) = \det \mathring{N}_{\zeta}(\lambda), \quad \lambda^2 < 0.$$

Observe that since all operators can be seen as acting on $L^2(\mathring{X})$ we have

$$\begin{aligned} e^{-t\Delta} - e^{-t\mathring{\Delta}} &= (e^{-t\Delta} - e^{-t\Delta^D}) - (e^{-t\mathring{\Delta}} - e^{-t\mathring{\Delta}^D}) + (e^{-t\Delta^D} - e^{-t\mathring{\Delta}^D}) \\ &= (e^{-t\Delta} - e^{-t\Delta^D}) - (e^{-t\mathring{\Delta}} - e^{-t\mathring{\Delta}^D}) + (e^{-t\Delta^D_-} - e^{-t\mathring{\Delta}^D_-}), \end{aligned}$$

where for the last line we have used that $\Delta^D = \Delta^D_- \oplus \Delta^D_+$, $\mathring{\Delta}^D = \mathring{\Delta}^D_- \oplus \mathring{\Delta}^D_+$ since X_+ and \mathring{X}_+ are isometric.

It follows that we can take the trace of both sides and thus define the following relative zeta-function for $\Re s > 1$:

$$\zeta(s; \Delta - \lambda^2, \mathring{\Delta} - \lambda^2) = \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} e^{\lambda^2 t} \text{Tr}(e^{-t\Delta} - e^{-t\mathring{\Delta}}) dt, \quad \lambda^2 < 0.$$

Moreover, we obtain the relation

$$\begin{aligned} \zeta(s; \Delta - \lambda^2, \mathring{\Delta} - \lambda^2) &= \zeta(s; \Delta - \lambda^2, \Delta^D - \lambda^2) - \zeta(s; \mathring{\Delta} - \lambda^2, \mathring{\Delta}^D - \lambda^2) \\ &\quad + \zeta(s, \Delta^D_- - \lambda^2) - \zeta(s, \mathring{\Delta}^D_- - \lambda^2). \end{aligned}$$

All the functions continue meromorphically to the complex plane with no pole at 0. Passing to the determinant, we obtain

$$\frac{\det_\zeta(\Delta - \lambda^2, \Delta^D - \lambda^2)}{\det_\zeta(\mathring{\Delta} - \lambda^2, \mathring{\Delta}^D - \lambda^2)} = \frac{\det_\zeta(\Delta - \lambda^2, \mathring{\Delta} - \lambda^2) \det_\zeta(\mathring{\Delta}^D_- - \lambda^2)}{\det_\zeta(\Delta^D_- - \lambda^2)}.$$

Thus dividing (2.16) by (2.17) we obtain

$$(2.18) \quad \frac{\det_\zeta(\Delta - \lambda^2, \mathring{\Delta} - \lambda^2) \det_\zeta(\mathring{\Delta}^D_- - \lambda^2)}{\det_\zeta(\Delta^D_- - \lambda^2)} = \frac{\det_\zeta \mathcal{N}(\lambda)}{\det_\zeta \mathring{\mathcal{N}}(\lambda)}, \quad \lambda^2 < 0,$$

where $\mathring{\mathcal{N}}(\lambda)$ and $\mathring{\Delta}^D_-$ are moduli independent.

In order to take the limit $\lambda^2 \rightarrow 0-$ in (2.18), we will need the asymptotic behavior of all the ingredients in the latter equation. We start with $\det_\zeta \mathcal{N}(\lambda)$.

2.3. Asymptotic of $\det_\zeta \mathcal{N}(\lambda)$ as $|\lambda| \rightarrow 0+$, $\Im \lambda \geq 0$. In this section we follow [6], where a similar problem is studied. Our purpose here is to refine the asymptotic of $\det_\zeta \mathcal{N}(\lambda)$ from [6] as needed for the proof of our gluing formula.

First we need to understand the behavior of the internal and external Dirichlet-to-Neumann operators. Since the internal Dirichlet Laplacian Δ^D_- is positive, there is no problem in letting λ go to 0 in the definition of $\mathcal{N}_-(\lambda)$.

Concerning the external Dirichlet-to-Neumann operator, by separation of variables in each conical end, we see that in the case $\lambda = 0$ the exterior Dirichlet problem (2.8) has a unique solution of the form

$$(2.19) \quad u(r, \varphi; 0) = \sum_{n=-\infty}^\infty C_n \left(\frac{R}{r}\right)^{\nu_n} e^{in\varphi},$$

where $f = \sum C_n e^{in\varphi}$. We recall that $\nu_n = \frac{|n|}{kR}$ and $\psi_n(\phi) = (2\pi)^{-\frac{1}{2}} e^{in\phi}$. The external Dirichlet-to-Neumann $\mathcal{N}_+(0)$ is obtained by applying $-\partial_r$ to this solution. Clearly, $\{|n|/(kR^2), \psi_n\}_{n=-\infty}^\infty$ is a complete set of the eigenvalues and orthonormal eigenfunctions of the operator $\mathcal{N}_+(0)$.

Remark 1. Note that thanks to the special choice of the lower bound on y in (2.7) the eigenvalue $\mu_0(\lambda)$ of $\mathcal{N}_+(\lambda)$ corresponding to the constant eigenfunction ψ_0 does not depend on k . This will be important in our proof of the BFK gluing formula in the case $K \geq 1$ if $k_j \neq k_i$ for some $i, j = 1, \dots, K$.

It is convenient to present the argument in the case where $K = 1$ so that X has only one conical end. We will explain afterwards how the proof is modified for $K > 1$.

2.3.1. *The case $K = 1$.* In the series (2.19) only the terms with $\nu_n > 1$ are in $L^2(X_+)$. As a result, in a neighborhood of zero, properties of $\lambda \mapsto \mathcal{N}_+(\lambda)$ on the eigenspaces of $\mathcal{N}_+(0)$ corresponding to the eigenvalues $|n|/(kR^2) > 1/R$ and $|n|/(kR^2) \leq 1/R$ are essentially different. Consider the spectral projector $\mathbf{P} = \sum_{0 \leq n \leq kR} P_n$ of $\mathcal{N}_+(0)$ on the interval $[0, 1/R]$; here

$$P_0 = (\cdot, \psi_0)_{L^2(\Sigma)}; \quad P_n = (\cdot, \psi_n)_{L^2(\Sigma)}\psi_n + (\cdot, \psi_{-n})_{L^2(\Sigma)}\psi_{-n}.$$

Lemma 1 (See [6, Prop 4.5]). *We have*

$$\mathcal{N}_+(\lambda)(\text{Id} - \mathbf{P}) = \Psi(\lambda^2) + \mathcal{L}(\lambda^2),$$

where $\Psi(z)$ is an elliptic pseudodifferential operator of order 1 which is a holomorphic function of z in a neighborhood of zero and $\mathcal{L}(z)$ is an operator with a smooth integral kernel which is a C^1 function of z in a neighborhood of zero with $\Im z \geq 0$.

Recall that the eigenfunctions ψ_n of $\mathcal{N}_+(\lambda)$ do not depend on λ and we have

$$\mathcal{N}_+(\lambda)\mathbf{P}f = \sum_{0 \leq n \leq kR} \mu_n(\lambda)P_n,$$

where the μ_n have been defined in (2.10).

The eigenvalues of $\mathcal{N}_+(0)$ on $[0, 1/R]$ are the limits of $\mu_n(\lambda)$. As $|\lambda| \rightarrow 0+$, $\Im \lambda \geq 0$, the formula (2.10) and properties of the Hankel functions (see [1]) imply that

$$(2.20) \quad \mu_0(\lambda) = -\frac{1}{R \ln \lambda} \left(1 - \left(\ln \frac{R}{2} + \frac{\pi\gamma}{2} - i\frac{\pi}{2} \right) \frac{1}{\ln \lambda} + O\left(\frac{1}{(\ln \lambda)^2}\right) \right),$$

where γ is the Euler’s constant, and

$$(2.21) \quad \mu_n(\lambda) = |n|/(kR^2) + O(\lambda^\epsilon), \quad 0 < |n| < kR,$$

with some $\epsilon > 0$.

We show the following proposition.

Proposition 1. *Assume $(X, |df|^2)$ has only one conical end; then, for any R large enough we have, as $|\lambda| \rightarrow 0+$, $\Im \lambda \geq 0$,*

$$\det_\zeta \mathcal{N}(\lambda) = -\frac{1}{R \ln \lambda} \det_\zeta^* \mathcal{N}(0) \left(1 - \left(\ln \frac{R}{2} + \frac{\pi\gamma}{2} - i\frac{\pi}{2} \right) \frac{1}{\ln \lambda} + O\left(\frac{1}{(\ln \lambda)^2}\right) \right),$$

where $\det_\zeta \mathcal{N}(\lambda)$ is the zeta-regularized determinant of $\mathcal{N}(\lambda)$, and $\det_\zeta^* \mathcal{N}(0)$ is the zeta-regularized determinant of $\mathcal{N}(0)$ with zero eigenvalue excluded.

Proof. Due to the representation $L^2(\Sigma) = \ker P_0 \oplus \ker(\mathbf{P} - P_0) \oplus \ker(\text{Id} - \mathbf{P})$ we have

$$(2.22) \quad \mathcal{N}(\lambda) = \begin{pmatrix} \mathcal{N}_{0,0}(\lambda) & \mathcal{N}_{0,1}(\lambda) & \mathcal{N}_{0,2}(\lambda) \\ \mathcal{N}_{1,0}(\lambda) & \mathcal{N}_{1,1}(\lambda) & \mathcal{N}_{1,2}(\lambda) \\ \mathcal{N}_{2,0}(\lambda) & \mathcal{N}_{2,1}(\lambda) & \mathcal{N}_{2,2}(\lambda) \end{pmatrix},$$

where $\mathcal{N}_{i,j}(\lambda) = \mathcal{P}_i \mathcal{N}(\lambda) \mathcal{P}_j$ with $\mathcal{P}_0 = P_0$, $\mathcal{P}_1 = \mathbf{P} - P_0$, and $\mathcal{P}_2 = \text{Id} - \mathbf{P}$.

The operator $\mathcal{N}_{2,2}(0)$ is invertible and, therefore, $\det_\zeta \mathcal{N}_{2,2}(0) \neq 0$.

Note that $\mathcal{N}_{2,2}(\lambda) = \mathcal{P}_2 \mathcal{N}_-(\lambda) \mathcal{P}_2 + \mathcal{P}_2 \mathcal{N}_+(\lambda) \mathcal{P}_2$, where $\mathcal{N}_+(\lambda) \mathcal{P}_2 = \mathcal{N}_+(\lambda)(\text{Id} - \mathbf{P})$ is the same as in Lemma 1 and $\mathcal{N}_-(\lambda)$ is a holomorphic function of λ^2 in a small neighborhood of zero. This implies that

$$(2.23) \quad \det_\zeta \mathcal{N}_{2,2}(\lambda) - \det_\zeta \mathcal{N}_{2,2}(0) = o(1), \quad |\lambda| \rightarrow 0+, \Im \lambda \geq 0.$$

Thanks to Lemma 1 we also have

$$(2.24) \quad \begin{aligned} &\|\mathcal{N}_{2,2}(\lambda) - \mathcal{N}_{2,2}(0); \mathcal{B}(H^1(\Sigma), L^2(\Sigma))\| = O(\lambda^2), \\ &\|\partial_\lambda \mathcal{N}_{2,2}(\lambda) - \partial_\lambda \mathcal{N}_{2,2}(0); \mathcal{B}(H^1(\Sigma), L^2(\Sigma))\| = O(\lambda). \end{aligned}$$

In order to refine (2.23), we estimate the absolute value of $\partial_\lambda \ln \det_\zeta \mathcal{N}_{2,2}(\lambda)$. Since $\partial_\lambda \mathcal{N}_{2,2}(\lambda)$ and $\mathcal{N}_{2,2}^{-1}(\lambda)$ are pseudodifferential operators of order -1 , the operator

$$\mathcal{N}_{2,2}^{-1}(\lambda) \partial_\lambda \mathcal{N}_{2,2}(\lambda)$$

is in the trace class, and hence

$$(2.25) \quad \partial_\lambda \ln \det_\zeta \mathcal{N}_{2,2}(\lambda) = \text{Tr}\{\mathcal{N}_{2,2}^{-1}(\lambda) \partial_\lambda \mathcal{N}_{2,2}(\lambda)\};$$

see [4, 10]. The first estimate in (2.24) and the Neumann series for $\mathcal{N}_{2,2}^{-1}(\lambda)$ give

$$(2.26) \quad \begin{aligned} \mathcal{N}_{2,2}^{-1}(\lambda) &= (\text{Id} + L(\lambda))\mathcal{N}_{2,2}^{-1}(0), & \|L(\lambda); \mathcal{B}(H^1(\Sigma))\| &= O(\lambda^2); \\ \mathcal{N}_{2,2}^{-1}(\lambda) &= \mathcal{N}_{2,2}^{-1}(0)(\text{Id} + R(\lambda)), & \|R(\lambda); \mathcal{B}(L^2(\Sigma))\| &= O(\lambda^2). \end{aligned}$$

As a consequence of (2.25), (2.26), and (2.24) we get

$$\begin{aligned} |\partial_\lambda \ln \det_\zeta \mathcal{N}_{2,2}(\lambda)| &= |\text{Tr}\{\mathcal{N}_{2,2}^{-1}(\lambda) \partial_\lambda \mathcal{N}_{2,2}(\lambda)\}| \\ &\leq \|\mathcal{N}_{2,2}(\lambda)(\text{Id} + L(\lambda))\mathcal{N}_{2,2}^{-2}(0)(\text{Id} + R(\lambda))\partial_\lambda \mathcal{N}_{2,2}(\lambda)\|_1 \\ &\leq \|\mathcal{N}_{2,2}(\lambda); \mathcal{B}(H^1(\Sigma), L^2(\Sigma))\| \|\text{Id} + L(\lambda); \mathcal{B}(H^1(\Sigma))\| \|\mathcal{N}_{2,2}^{-2}(0)\|_1 \\ &\quad \times \|\text{Id} - R(\lambda); \mathcal{B}(L^2(\Sigma))\| \|\partial_\lambda \mathcal{N}_{2,2}(\lambda); \mathcal{B}(H^1(\Sigma), L^2(\Sigma))\| = O(1), \end{aligned}$$

where $\|\cdot\|_1$ is the trace norm. This together with (2.23) implies $|\partial_\lambda \det_\zeta \mathcal{N}_{2,2}(\lambda)| = O(1)$. Now, as a refinement of (2.23), we obtain

$$\det_\zeta \mathcal{N}_{2,2}(\lambda) - \det_\zeta \mathcal{N}_{2,2}(0) = O(\lambda).$$

This together with (2.22) implies

$$(2.27) \quad \begin{aligned} \det_\zeta \mathcal{N}(\lambda) &= \det_{\text{Fr}} \begin{pmatrix} \mathcal{N}_{0,0}(\lambda) & \mathcal{N}_{0,1}(\lambda) & \mathcal{N}_{0,2}(\lambda)\mathcal{N}_{2,2}(\lambda)^{-1} \\ \mathcal{N}_{1,0}(\lambda) & \mathcal{N}_{1,1}(\lambda) & \mathcal{N}_{1,2}(\lambda)\mathcal{N}_{2,2}(\lambda)^{-1} \\ \mathcal{N}_{2,0}(\lambda) & \mathcal{N}_{2,1}(\lambda) & \text{Id} \end{pmatrix} \det_\zeta \begin{pmatrix} \text{Id} & 0 & 0 \\ 0 & \text{Id} & 0 \\ 0 & 0 & \mathcal{N}_{2,2}(\lambda) \end{pmatrix} \\ &= \det_{\text{Fr}} \begin{pmatrix} \mathcal{N}_{0,0}(\lambda) & \mathcal{N}_{0,1}(\lambda) & \mathcal{N}_{0,2}(\lambda)\mathcal{N}_{2,2}(\lambda)^{-1} \\ \mathcal{N}_{1,0}(\lambda) & \mathcal{N}_{1,1}(\lambda) & \mathcal{N}_{1,2}(\lambda)\mathcal{N}_{2,2}(\lambda)^{-1} \\ \mathcal{N}_{2,0}(\lambda) & \mathcal{N}_{2,1}(\lambda) & \text{Id} \end{pmatrix} (\det_\zeta \mathcal{N}_{2,2}(0))(1 + O(\lambda)); \end{aligned}$$

see [28] for the first equality. On the next step we rely on the estimate

$$(2.28) \quad |\det_{\text{Fr}}(\text{Id} + A) - \det_{\text{Fr}}(\text{Id} + B)| \leq \|A - B\|_1 e^{\|A\|_1 + \|B\|_1 + 1},$$

see [37, f-la (3.7), and the references therein], for

$$\begin{aligned} \text{Id} + A &= \begin{pmatrix} \mathcal{N}_{0,0}(\lambda) & 0 & 0 \\ 0 & \mathcal{N}_{1,1}(0) & \mathcal{N}_{1,2}(0)\mathcal{N}_{2,2}^{-1}(0) \\ 0 & \mathcal{N}_{2,1}(0) & \text{Id} \end{pmatrix}, \\ \text{Id} + B &= \begin{pmatrix} \mathcal{N}_{0,0}(\lambda) & \mathcal{N}_{0,1}(\lambda) & \mathcal{N}_{0,2}(\lambda)\mathcal{N}_{2,2}(\lambda)^{-1} \\ \mathcal{N}_{1,0}(\lambda) & \mathcal{N}_{1,1}(\lambda) & \mathcal{N}_{1,2}(\lambda)\mathcal{N}_{2,2}(\lambda)^{-1} \\ \mathcal{N}_{2,0}(\lambda) & \mathcal{N}_{2,1}(\lambda) & \text{Id} \end{pmatrix}. \end{aligned}$$

Since $\mathcal{N}_-(0)$ is a self-adjoint operator in $L^2(\Sigma)$ and $\ker \mathcal{N}_-(0) = \{c \in \mathbb{C}\}$, we have $\mathcal{N}_-(0)P_0 = P_0\mathcal{N}_-(0) = 0$. Then thanks to

$$P_i(\mathcal{N}(\lambda) - \mathcal{N}(0))P_j = \delta_{ij}(\mu_j(\lambda) - \mu_j(0)) + P_i(\mathcal{N}_-(\lambda) - \mathcal{N}_-(0))P_j, \quad i, j \in [0, 1/R],$$

where δ_{ij} is the Kronecker delta function, and

$$P_j(\mathcal{N}(\lambda) - \mathcal{N}(0))(\text{Id} - \mathbf{P}) = P_j(\mathcal{N}_-(\lambda) - \mathcal{N}_-(0))(\text{Id} - \mathbf{P}), \quad j \in [0, 1/R],$$

together with (2.21) and (2.26), we obtain $\|A - B\|_1 = O(\lambda^\epsilon)$ with some $\epsilon > 0$. From (2.27) and (2.28) we get

$$(2.29) \quad \det_\zeta \mathcal{N}(\lambda) = \det_{\text{Fr}} \begin{pmatrix} \mathcal{N}_{0,0}(\lambda) & 0 & 0 \\ 0 & \mathcal{N}_{1,1}(0) & \mathcal{N}_{1,2}(0)\mathcal{N}_{2,2}(0)^{-1} \\ 0 & \mathcal{N}_{2,1}(0) & \text{Id} \end{pmatrix} \det_\zeta \mathcal{N}_{2,2}(0)(1 + O(\lambda^\epsilon)).$$

It remains to note that

$$\begin{aligned} \mathcal{N}_{0,0}(\lambda) &= (\mu_0(\lambda) + P_0(\mathcal{N}_-(\lambda) - \mathcal{N}_-(0)))P_0 = (\mu_0(\lambda) + O(\lambda^2))P_0, \\ \det_\zeta^* \mathcal{N}(0) &= \det_{\text{Fr}} \begin{pmatrix} \mathcal{N}_{1,1}(0) & \mathcal{N}_{1,2}(0)\mathcal{N}_{2,2}(0)^{-1} \\ \mathcal{N}_{2,1}(0) & \text{Id} \end{pmatrix} \det_\zeta \mathcal{N}_{2,2}(0). \end{aligned}$$

This together with (2.29) and (2.20) completes the proof. □

Corollary 1. *The spectral shift function ξ in (2.14) satisfies*

$$(2.30) \quad \xi(\lambda; \Delta, \Delta^D) = (\ln \lambda^2)^{-1} + O((\ln \lambda)^{-2}), \quad \lambda \rightarrow 0+.$$

Proof. By [5, Theorem 3.5] we have $\xi(\lambda) = \pi^{-1} \text{Arg} \det \mathcal{N}(\sqrt{\lambda^2 + i0})$ as $\lambda^2 \rightarrow 0+$, where $\text{Arg} z \in (-\pi, \pi]$, and $\xi(\lambda) = 0$ if $\lambda^2 < 0$. Calculation of the argument in the asymptotic obtained in Proposition 1 gives (2.30). □

2.3.2. The case $K > 1$. Let us outline the changes in Proposition 1 and Corollary 1 needed in the case $K > 1$. Now we have $\mathcal{N}_+(\lambda) = \bigoplus_{j=1}^K \mathcal{N}_+^{(j)}(\lambda)$, where each $\mathcal{N}_+^{(j)}(\lambda)$ is defined on the circle $\{y \in \mathbb{C}_j : |y| = R^{1/k_j}\}$ as in (2.9). The first eigenvalue of $\mathcal{N}_+^{(j)}(\lambda)$ is $\mu_0(\lambda)$ and the corresponding eigenspace consists of constant functions on the circle. As a consequence, in the estimate (2.29) the eigenvectors in \mathcal{N}_+ with eigenvalue $\mu_0(\lambda)$ contribute at the order $O(\frac{1}{\ln \lambda})$ instead of $O(\lambda^\epsilon)$ and this is not good enough for our purpose.

We thus introduce \mathbf{P}_0 the orthogonal projection onto the eigenspace of $\mathcal{N}_+(\lambda)$ corresponding to $\mu_0(\lambda)$ (note that \mathbf{P}_0 does not depend on λ , and that $\text{rank } \mathbf{P}_0 = K$). Observe that we have $\ker \mathcal{N}(0) \subset \ker(\text{Id} - \mathbf{P}_0)$. We repeat the argument of Proposition 1, where \mathcal{P}_0 is now the orthogonal projection onto $\ker \mathcal{N}(0) = \{c \in \mathbb{C}\}$, $\mathcal{P}_1 = (\text{Id} - \mathcal{P}_0)\mathbf{P}$ where \mathbf{P} is the spectral projection of $\mathcal{N}_+(0)$ on the interval $[0, 1/R]$, and $\mathcal{P}_2 = (\text{Id} - \mathbf{P})$. Clearly, $\mathcal{N}_+(\lambda)\mathcal{P}_0 = \mu_0(\lambda)\mathcal{P}_0$ and $\mathcal{N}_+(\lambda)(\text{Id} - \mathcal{P}_0) = (\text{Id} - \mathcal{P}_0)\mathcal{N}_+(\lambda)$.

The same argument as in the case $K = 1$ leads to

$$\begin{aligned} \det_\zeta \mathcal{N}(\lambda) &= -\frac{1}{R \ln \lambda} \det_{\text{Fr}} \begin{pmatrix} \mathcal{N}_{1,1}(0) + \mu_0(\lambda)\mathbf{P}_0(\text{Id} - \mathcal{P}_0) & \mathcal{N}_{1,2}(0)\mathcal{N}_{2,2}(0)^{-1} \\ \mathcal{N}_{2,1}(0) & \text{Id} \end{pmatrix} \\ &\quad \times \det_\zeta \mathcal{N}_{2,2}(0) \left(1 - \left(\ln \frac{R}{2} + \frac{\pi\gamma}{2} - i\frac{\pi}{2} \right) \frac{1}{\ln \lambda} + O\left(\frac{1}{(\ln \lambda)^2}\right) \right). \end{aligned}$$

(Note that in the case $K = 1$ we have $\mathbf{P}_0 = \mathcal{P}_0$ and the term $\mu_0(\lambda)\mathbf{P}_0(\text{Id} - \mathcal{P}_0)$ does not appear.) This together with (2.20) and (2.28) gives

$$\begin{aligned} \det_\zeta \mathcal{N}(\lambda) &= -\frac{1}{R \ln \lambda} \det_{\text{Fr}} \begin{pmatrix} \mathcal{N}_{1,1}(0) - \frac{1}{R \ln \lambda} \mathbf{P}_0(\text{Id} - \mathcal{P}_0) & \mathcal{N}_{1,2}(0)\mathcal{N}_{2,2}(0)^{-1} \\ \mathcal{N}_{2,1}(0) & \text{Id} \end{pmatrix} \\ &\quad \times \det_\zeta \mathcal{N}_{2,2}(0) \left(1 - \left(\ln \frac{R}{2} + \frac{\pi\gamma}{2} - i\frac{\pi}{2} \right) \frac{1}{\ln \lambda} + O\left(\frac{1}{(\ln \lambda)^2}\right) \right). \end{aligned}$$

Here the Fredholm determinant is a holomorphic function of the parameter $\tau := \frac{1}{R \ln \lambda}$. We have

$$(2.31) \quad \det_{\zeta} \mathcal{N}(\lambda) = -\frac{1}{R \ln \lambda} \det_{\zeta}^* \mathcal{N}(0) \left(1 - \left(C + \ln \frac{R}{2} + \frac{\pi\gamma}{2} - i\frac{\pi}{2} \right) \frac{1}{\ln \lambda} + O\left(\frac{1}{(\ln \lambda)^2}\right) \right)$$

with some constant $C = C(R)$.

We observe that C must be real since $\det_{\zeta} \mathcal{N}(\lambda)$ is positive for $\lambda \in \mathbb{C}$, $\text{Arg } \lambda = \pi/2$ (as $\mathcal{N}(\lambda)$ is a positive self-adjoint operator for those values of λ). Thus C does not influence the calculation of the argument in the asymptotic of $\det_{\zeta} \mathcal{N}(\lambda)$ and Corollary 1 remains valid for $K > 1$. We will use Corollary 1 to define a relative determinant of $(\Delta, \mathring{\Delta})$ at $\lambda = 0$.

2.4. The relative determinant and the gluing formula at $\lambda = 0$. In this section we prove the following theorem.

Theorem 4. *The gluing formula*

$$\det_{\zeta}(\Delta, \mathring{\Delta}) = C \det_{\zeta}^* \mathcal{N} \cdot \det_{\zeta} \Delta_{-}^D$$

is valid, where \mathcal{N} , Δ_{-}^D depend on R . The constant C depends on R but not on the moduli parameters z_1, \dots, z_M .

Observe that this theorem first requires a definition for the left-hand side of the equality. Once this is done, we will let λ go to zero in (2.18) and study the limit of both sides.

As before the case $K = 1$ is simpler than the general one. We will present the proof for this case first. The case $K > 1$ is more technically involved but the arguments we need can be adapted from [14].

2.4.1. The case $K = 1$. In this case, the definition of $\det_{\zeta}(\Delta, \mathring{\Delta})$ is rather straightforward, since, for $K = 1$, a conventional regularization (see, e.g., [16] and the references therein) for the relative zeta-function makes sense. Indeed, the first integral in the representation

$$(2.32) \quad \zeta(s; \Delta, \mathring{\Delta}) = \left(\int_0^1 + \int_1^{\infty} \right) \frac{t^{s-1}}{\Gamma(s)} \text{Tr}(e^{-t\Delta} - e^{-t\mathring{\Delta}}) dt$$

defines an analytic function in $\Re s > 1$ that has a meromorphic continuation to \mathbb{C} with no pole at zero by the usual argument based on short time heat trace asymptotic

$$(2.33) \quad \text{Tr}(e^{-t\Delta} - e^{-t\mathring{\Delta}}) \sim \sum_{j \geq -2} a_j t^{j/2}, \quad t \rightarrow 0+.$$

For the second integral we need the long time heat trace behavior given by the following lemma.

Lemma 2. *Assume that $K = 1$. Then*

$$\text{Tr}(e^{-t\Delta} - e^{-t\mathring{\Delta}}) = O((\ln t)^{-2}) \text{ as } t \rightarrow +\infty.$$

Proof. Since $\Delta_+^D \equiv \mathring{\Delta}_+^D$ and the operators $\Delta_-^D, \mathring{\Delta}_-^D$ are Dirichlet Laplacians on compact manifolds, we have

$$\begin{aligned}
 (2.34) \quad & \text{Tr}(e^{-t\Delta} - e^{-t\mathring{\Delta}}) \\
 &= \text{Tr}(e^{-t\Delta} - e^{-t\Delta_-^D \oplus \Delta_+^D}) - \text{Tr}(e^{-t\mathring{\Delta}} - e^{-t\mathring{\Delta}_-^D \oplus \mathring{\Delta}_+^D}) + \text{Tr} e^{-t\Delta_-^D} - \text{Tr} e^{-t\mathring{\Delta}_-^D} \\
 &= -t \int_0^\infty e^{-t\lambda^2} (\xi(\lambda; \Delta, \Delta^D) - \xi(\lambda; \mathring{\Delta}, \mathring{\Delta}^D)) 2\lambda d\lambda + O(e^{-t\delta}), \quad t \rightarrow +\infty,
 \end{aligned}$$

where $\delta > 0$ is the smallest eigenvalue in the spectra of Δ_-^D and $\mathring{\Delta}_-^D$. (In (2.34) we also used (2.14) for Δ and $\mathring{\Delta}$.) As a consequence of Corollary 1 (which is also valid for $\mathring{\xi}$ in the case $K = 1$) we have

$$\xi(\lambda; \Delta, \Delta^D) - \xi(\lambda, \mathring{\Delta}, \mathring{\Delta}^D) = O((\ln \lambda)^{-2}), \quad \lambda \rightarrow 0+.$$

This together with (2.34) implies the assertion; see, e.g., [17, Theorem 1.7] for details. □

As a consequence, the second integral in (2.32) defines a holomorphic function in $\Re s < 0$ that has a continuous in $\Re s \leq 0$ derivative. Thus $\zeta(s; \Delta, \mathring{\Delta})$ is a meromorphic function in $\Re s < 0$ and $\zeta'(s; \Delta, \mathring{\Delta})$ tends to a certain limit $\zeta'(0; \Delta, \mathring{\Delta})$ as $s \rightarrow 0-$. The relative zeta-regularized determinant is defined to be

$$(2.35) \quad \det_\zeta(\Delta, \mathring{\Delta}) = e^{-\zeta'(0; \Delta, \mathring{\Delta})}.$$

We now prove the gluing formula in the case $K = 1$. First observe that by Proposition 1 (applied also to $\mathring{N}(\lambda)$) we have

$$(2.36) \quad \frac{\det_\zeta N(\lambda)}{\det_\zeta \mathring{N}(\lambda)} \rightarrow \frac{\det_\zeta^* N(0)}{\det_\zeta^* \mathring{N}(0)} \text{ as } \lambda^2 \rightarrow 0-, \quad K = 1.$$

The limit $\lambda \rightarrow 0$ is then addressed by the

Proposition 2. *In the case $K = 1$ we have*

$$\det_\zeta(\Delta - \lambda^2, \mathring{\Delta} - \lambda^2) \rightarrow \det_\zeta(\Delta, \mathring{\Delta}) \text{ as } \lambda^2 \rightarrow 0-,$$

where the determinant $\det_\zeta(\Delta, \mathring{\Delta})$ has been defined in (2.35).

Proof. Let us write the relative zeta-function in the form

$$\zeta(s; \Delta - \lambda^2, \mathring{\Delta} - \lambda^2) = \left(\int_0^1 + \int_1^\infty \right) \frac{t^{s-1} e^{t\lambda^2}}{\Gamma(s)} \text{Tr}(e^{-t\Delta} - e^{-t\mathring{\Delta}}) dt.$$

Thanks to (2.15) the first integral converges for $\Re s > 1$ uniformly in $\lambda \leq 0$ and has a meromorphic continuation to \mathbb{C} with no pole at zero (by the usual argument based on the short time heat trace asymptotic (2.33)). Due to Lemma 2 the second integral defines a holomorphic in $\Re s < 0$ and continuous in $\lambda^2 \leq 0$ and $\Re s \leq 0$ function. Moreover, as $1/\Gamma(s)$ has a first order zero at $s = 0$, Lemma 2 also implies that the first derivative with respect to s of the second integral is also continuous in $\lambda^2 \leq 0$ and $\Re s \leq 0$. Thus we obtain

$$\zeta'(0; \Delta - \lambda^2, \mathring{\Delta} - \lambda^2) \rightarrow \zeta'(0; \Delta, \mathring{\Delta}), \quad \lambda^2 \rightarrow 0-,$$

where $\zeta'(0; \Delta, \mathring{\Delta})$ is defined using (2.32). □

Proof of Theorem 4 in the case $K = 1$. We pass to the limit as $\lambda^2 \rightarrow 0-$ in (2.18). Since Δ_-^D is positive, we have $\det_\zeta(\Delta_-^D - \lambda^2) \rightarrow \det_\zeta \Delta_-^D$ as $\lambda^2 \rightarrow 0-$, and the same is true for $\mathring{\Delta}_-^D$. Thanks to (2.36) and Proposition 2 we obtain

$$\frac{\det_\zeta(\Delta, \mathring{\Delta}) \det_\zeta \mathring{\Delta}_-^D}{\det_\zeta \Delta_-^D} = \frac{\det_\zeta^* \mathcal{N}(0)}{\det_\zeta^* \mathring{\mathcal{N}}(0)},$$

which proves Theorem 4, where $\mathcal{N}(0)$ is denoted by \mathcal{N} and the constant

$$C = (\det_\zeta \mathring{\Delta}_-^D \det_\zeta^* \mathring{\mathcal{N}}(0))^{-1}$$

is moduli independent. □

2.4.2. *The case $K > 1$.* In the case $K > 1$ we have $\mathring{\mathcal{N}}(\lambda) = \bigoplus_{j=1}^K \mathring{\mathcal{N}}^{(j)}(\lambda)$, where $\mathring{\mathcal{N}}^{(j)}(\lambda)$ is the Neumann jump operator on the circle $\{y \in \mathbb{C}_j : |y| = R^{1/k_j}\}$ located on the infinite cone $(\mathbb{C}_j, |dy^{k_j}|^2)$. We have

$$\det_\zeta \mathring{\mathcal{N}}(\lambda) = \prod_{j=1}^K \det_\zeta \mathring{\mathcal{N}}^{(j)}(\lambda), \quad \det_\zeta^* \mathring{\mathcal{N}}(0) = \prod_{j=1}^K \det_\zeta^* \mathring{\mathcal{N}}^{(j)}(0).$$

We apply Proposition 1 to each $\det_\zeta \mathring{\mathcal{N}}^{(j)}(\lambda)$, $j = 1, \dots, K$ and get (2.37)

$$\det_\zeta \mathring{\mathcal{N}}(\lambda) = (-R \ln \lambda)^{-K} \det_\zeta^* \mathcal{N}(0) \left(1 - \left(\ln \frac{R}{2} + \frac{\pi\gamma}{2} - i\frac{\pi}{2} \right) \frac{K}{\ln \lambda} + O\left(\frac{1}{(\ln \lambda)^2}\right) \right),$$

as $|\lambda| \rightarrow 0+$, $\Im \lambda \geq 0$.

Thanks to the relation $\xi(\lambda; \mathring{\Delta}, \mathring{\Delta}^D) = \pi^{-1} \text{Arg} \det \mathring{\mathcal{N}}(\sqrt{\lambda^2 + i0})$ as $\lambda^2 \rightarrow 0+$, calculation of the argument in (2.37) leads to

$$\xi(\lambda; \mathring{\Delta}, \mathring{\Delta}^D) = K(\ln \lambda^2)^{-1} + O((\ln \lambda)^{-2}), \quad \lambda \rightarrow 0+,$$

where $\xi(\cdot; \mathring{\Delta}, \mathring{\Delta}^D) \in L^1(\mathbb{R}_+, (1 + \lambda^2)^{-2} d\lambda^2)$ is the spectral shift function satisfying

$$\text{Tr}((\mathring{\Delta} + 1)^{-1} - (\mathring{\Delta}^D + 1)^{-1}) = - \int_0^\infty \xi(\lambda; \mathring{\Delta}, \mathring{\Delta}^D) (1 + \lambda^2)^{-2} d\lambda^2;$$

cf. Corollary 1. This together with Corollary 1 gives

$$(2.38) \quad \xi(\lambda, \Delta, \Delta^D) - \xi(\lambda, \mathring{\Delta}, \mathring{\Delta}^D) = -(K - 1)(\ln \lambda^2)^{-1} + O((\ln \lambda)^{-2}), \quad \lambda \rightarrow 0+.$$

Besides, Proposition 1 together with (2.37) implies that

$$(2.39) \quad \left(\ln \frac{i}{\lambda} \right)^{1-K} \frac{\det_\zeta \mathcal{N}(\lambda)}{\det_\zeta \mathring{\mathcal{N}}(\lambda)} \rightarrow R^{K-1} \frac{\det_\zeta^* \mathcal{N}(0)}{\det_\zeta^* \mathring{\mathcal{N}}(0)} \text{ as } \lambda^2 \rightarrow 0-, \quad K \geq 1.$$

Recall that for $\lambda^2 < 0$ the relative zeta-function is defined as the meromorphic continuation of

$$(2.40) \quad \zeta(s; \Delta - \lambda^2, \mathring{\Delta} - \lambda^2) = \int_0^\infty \frac{t^{s-1} e^{t\lambda^2}}{\Gamma(s)} \text{Tr}(e^{-t\Delta} - e^{-t\mathring{\Delta}}) dt$$

from $\Re s > 1$.

We have

$$\begin{aligned} & \text{Tr}(e^{-t\Delta} - e^{-t\mathring{\Delta}}) \\ &= \text{Tr}(e^{-t\Delta} - e^{-t\Delta^D \oplus \Delta_+^D}) - \text{Tr}(e^{-t\mathring{\Delta}} - e^{-t\mathring{\Delta}^D \oplus \mathring{\Delta}_+^D}) + \text{Tr} e^{-t\Delta^D} - \text{Tr} e^{-t\mathring{\Delta}^D}. \end{aligned}$$

(Now the short time asymptotic (2.33) is a consequence of (2.15) and similar short time asymptotics for $\text{Tr}(e^{-t\mathring{\Delta}} - e^{-t\mathring{\Delta}^D})$, $\text{Tr} e^{-t\Delta^D}$, and $\text{Tr} e^{-t\mathring{\Delta}^D}$.) Let

$$N(\lambda) = \sum_{j:\lambda_j^2 \leq \lambda^2} \dim \ker(\Delta_-^D - \lambda_j^2),$$

where λ_j^2 are the eigenvalues of Δ_-^D , be the counting function of Δ_-^D . Similarly, let $\mathring{N}(\lambda)$ be the counting function of $\mathring{\Delta}^D$. Then

$$\text{Tr}(e^{-t\Delta^D} - e^{-t\mathring{\Delta}^D}) = t \int_0^\infty e^{-t\lambda^2} (N(\lambda) - \mathring{N}(\lambda)) 2\lambda d\lambda$$

and

$$\xi(\lambda; \Delta; \mathring{\Delta}) = \xi(\lambda; \Delta, \Delta^D) - \xi(\lambda; \mathring{\Delta}, \mathring{\Delta}^D) - N(\lambda) + \mathring{N}(\lambda)$$

is the spectral shift function for the pair $(\Delta, \mathring{\Delta})$ such that

$$\text{Tr}(e^{-t\Delta} - e^{-t\mathring{\Delta}}) = -t \int_0^\infty e^{-t\lambda^2} \xi(\lambda; \Delta, \mathring{\Delta}) 2\lambda d\lambda.$$

Since the operators Δ_-^D and $\mathring{\Delta}^D$ are positive, from (2.38) it follows the asymptotic

$$(2.41) \quad \xi(\lambda; \Delta, \mathring{\Delta}) = -(K - 1)(\ln \lambda^2)^{-1} + O((\ln \lambda)^{-2}), \quad \lambda \rightarrow 0 +.$$

Introduce a cut-off function $\chi \in C^\infty(\mathbb{R})$ such that $\chi(\mu) = 1$ for $\mu < 1/2$ and $\chi(\mu) = 0$ for $\mu > 3/4$. Following the scheme in [14] we write

$$\text{Tr}(e^{-t\Delta} - e^{-t\mathring{\Delta}}) = e_1(t) + e_2(t),$$

where

$$e_1(t) = -t \int_0^\infty e^{-t\mu^2} \chi(\mu) \xi(\mu; \Delta, \mathring{\Delta}) 2\mu d\mu,$$

$$e_2(t) = -t \int_0^\infty e^{-t\mu^2} (1 - \chi(\mu)) \xi(\mu; \Delta, \mathring{\Delta}) 2\mu d\mu;$$

cf. (2.14). Note that e_2 is exponentially decreasing as $t \rightarrow +\infty$. Thanks to the short time asymptotic (2.33) and smoothness of e_1 at $t = 0$, we see that $e_2(t)$ has a short time asymptotic of the same form. Therefore, for $\lambda^2 \leq 0$ the holomorphic in $\Re s > 1$ zeta-function

$$\zeta_2(s; \lambda^2) = \int_0^\infty \frac{t^{s-1} e^{t\lambda^2}}{\Gamma(s)} e_2(t) dt$$

continues as a meromorphic function to \mathbb{C} with no pole at zero and $\zeta_2'(0, \lambda^2) \rightarrow \zeta_2'(0, 0)$ as $\lambda^2 \rightarrow 0-$ by the usual argument.

We are now in position to define the regularized determinant $\det_\zeta(\Delta, \mathring{\Delta})$. We start from $\zeta(s; \Delta, \mathring{\Delta}) = \zeta_1(s; 0) + \zeta_2(s; 0)$, where $\zeta_1(s; 0)$ is defined by

$$(2.42) \quad \zeta_1(s; 0) = \int_0^\infty \frac{t^{s-1}}{\Gamma(s)} e_1(t) dt = s \int_0^\infty (-\mu^2)^{-s-1} \chi(\mu) \xi(\mu; \Delta, \mathring{\Delta}) 2\mu d\mu.$$

From this expression and (2.38) one can easily see that $\zeta_1(s; 0)$ is a holomorphic function in $\Re s < 0$, and we already know that $\zeta_2(s; 0)$ is a meromorphic function of $s \in \mathbb{C}$ with no pole at zero.

The asymptotic behavior of $\zeta(s; 0)$ near $s = 0$ is given by the following proposition.

Proposition 3. *Set*

$$Z'_0 := \zeta'_2(0; 0) + (K - 1) \lim_{\delta \rightarrow 0^+} \left(\int_{\delta}^{\infty} \frac{\chi(\mu) d\mu}{\mu \ln \mu} + \ln \ln \frac{1}{\delta} \right) - 2 \int_0^{\infty} \mu^{-1} \chi(\mu) (\xi(\mu; \Delta, \mathring{\Delta}) - (K - 1)(\ln \mu^2)^{-1}) d\mu + (K - 1)(\gamma + \ln 2).$$

When $s \rightarrow 0^-$ we have

$$\zeta(s; \Delta, \mathring{\Delta}) = \zeta_2(0; 0) + s(K - 1) \ln(-s) + sZ'_0 + o(s).$$

This proposition gives way to the following definition of the regularized relative determinant of Δ in case $K > 1$.

Definition 4.

$$\det_{\zeta}(\Delta, \mathring{\Delta}) := e^{-Z'_0}.$$

Proof. Since $\zeta_2(s; 0)$ is a meromorphic function of s with no pole at zero, as $s \rightarrow 0^-$ we have

$$\zeta(s; \Delta, \mathring{\Delta}) = \zeta_1(s; 0) + \zeta_2(0; 0) + s\zeta'_2(0; 0) + O(s^2).$$

It remains to study the behavior of

$$\zeta_1(s; 0) = s \int_0^{\infty} (-\mu^2)^{-s-1} \chi(\mu) \xi(\mu; \Delta, \mathring{\Delta}) 2\mu d\mu$$

as $s \rightarrow 0^-$. We represent the last integral as a sum of two integrals. Due to (2.41) the first integral

$$s \int_0^{\infty} (-\mu^2)^{-s-1} \chi(\mu) (\xi(\mu; \Delta, \mathring{\Delta}) + (K - 1)(\ln \mu^2)^{-1}) 2\mu d\mu$$

converges uniformly in $s \leq 0$ and thus gives the contribution

$$-s \int_0^{\infty} 2\mu^{-1} \chi(\mu) (\xi(\mu; \Delta, \mathring{\Delta}) + (K - 1)(\ln \mu^2)^{-1}) d\mu$$

into the expansion of $\zeta(s; \Delta, \mathring{\Delta})$. For the second integral we have

$$s(1 - K) \int_0^{\infty} (-\mu^2)^{-s-1} \chi(\mu) (\ln \mu^2)^{-1} 2\mu d\mu = s(1 - K) \left(-\ln(-s) + \gamma + \ln 2 - \lim_{\delta \rightarrow 0^+} \left(\int_{\delta}^{\infty} \frac{\chi(\mu) d\mu}{\mu \ln \mu} + \ln \ln \frac{1}{\delta} \right) + o(1) \right);$$

see [14, p. 987]. □

The proof of the gluing formula will also require that we understand the limit when λ goes to 0. At this stage we have

$$\zeta(s; \Delta - \lambda^2, \mathring{\Delta} - \lambda^2) = \zeta_1(s; \lambda^2) + \zeta_2(s; \lambda^2),$$

where only properties of the zeta-function

$$(2.43) \quad \zeta_1(s; \lambda^2) = \int_0^{\infty} \frac{t^{s-1} e^{t\lambda^2}}{\Gamma(s)} e_1(t) dt = s \int_0^{\infty} (\lambda^2 - \mu^2)^{-s-1} \chi(\mu) \xi(\lambda; \Delta, \mathring{\Delta}) 2\mu d\mu$$

remain unknown. Notice that the last integrand is compactly supported and, therefore, the integral converges uniformly near $s = 0$ for fixed $\lambda^2 < 0$ due to (2.41). We get

$$(2.44) \quad \zeta'_1(0; \lambda^2) = \int_0^{\infty} (\lambda^2 - \mu^2)^{-1} \chi(\mu) \xi(\mu; \Delta, \mathring{\Delta}) 2\mu d\mu.$$

Proposition 4. *As $\lambda^2 \rightarrow 0-$ we have*

$$\begin{aligned} \zeta'(0; \Delta - \lambda^2, \mathring{\Delta} - \lambda^2) &= \ln \left(\ln \frac{i}{\lambda} \right)^{1-K} + (K-1) \lim_{\delta \rightarrow 0+} \left(\int_{\delta}^{\infty} \frac{\chi(\mu) d\mu}{\mu \ln \mu} + \ln \ln \frac{1}{\delta} \right) \\ &\quad - 2 \int_0^{\infty} \mu^{-1} \chi(\mu) \left(\xi(\mu; \Delta, \mathring{\Delta}) + (K-1)(\ln \mu^2)^{-1} \right) d\mu + \zeta'_2(0; 0) + o(1). \end{aligned}$$

Proof. We only need to study the behavior of $\zeta'(0; \lambda^2)$ in (2.44) as $\lambda^2 \rightarrow 0-$. Thanks to (2.41) the integral

$$\int_0^{\infty} (\lambda^2 - \mu^2)^{-1} \chi(\mu) \left(\xi(\mu; \Delta, \mathring{\Delta}) + (K-1)(\ln \mu^2)^{-1} \right) 2\mu d\mu$$

converges uniformly in $\lambda^2 \leq 0$ and thus tends to

$$- \int_0^{\infty} 2\mu^{-1} \chi(\mu) \left(\xi(\mu; \Delta, \mathring{\Delta}) + (K-1)(\ln \mu^2)^{-1} \right) d\mu$$

as $\lambda^2 \rightarrow 0-$. It remains to note that

$$\int_0^{\infty} (\lambda^2 - \mu^2)^{-1} \chi(\mu) (\ln \mu^2)^{-1} 2\mu d\mu = \ln \ln \frac{i}{\lambda} - \lim_{\delta \rightarrow 0+} \left(\int_{\delta}^{\infty} \frac{\chi(\mu) d\mu}{\mu \ln \mu} + \ln \ln \frac{1}{\delta} \right) + o(1);$$

as $\lambda^2 \rightarrow 0-$; see [14, p. 12 and appendix]. □

Note that by definition of $\det_{\zeta}(\Delta, \mathring{\Delta}) = e^{-Z'_0}$ we have

$$\zeta'(0; \Delta - \lambda^2, \mathring{\Delta} - \lambda^2) = \ln \left(\ln \frac{i}{\lambda} \right)^{1-K} - (-Z'_0 - (K-1)(\gamma + \ln 2)) + o(1), \quad \lambda^2 \rightarrow 0-;$$

see Proposition 4.

Proof of Theorem 4 in the general case. From Propositions 4 and 3 we immediately get

$$(2.45) \quad \left(\ln \frac{i}{\lambda} \right)^{1-K} \det_{\zeta}(\Delta - \lambda^2; \mathring{\Delta} - \lambda^2) \rightarrow e^{(1-K)(\gamma + \ln 2)} \det_{\zeta}^*(\Delta, \mathring{\Delta}), \quad \lambda^2 \rightarrow 0-.$$

We pass in (2.18) to the limit as $\lambda^2 \rightarrow 0-$. Taking into account (2.39) and (2.45) we obtain

$$\frac{\det_{\zeta}(\Delta, \mathring{\Delta}) \det_{\zeta} \mathring{\Delta}^D}{\det_{\zeta} \Delta^D} = (Re^{\gamma + \ln 2})^{K-1} \frac{\det_{\zeta}^* \mathcal{N}(0)}{\det_{\zeta}^* \mathring{\mathcal{N}}(0)}.$$

This proves Theorem 4, where $\mathcal{N} \equiv \mathcal{N}(0)$ and the constant

$$C = (Re^{\gamma + \ln 2})^{K-1} / (\det_{\zeta} \mathring{\Delta}^D \det_{\zeta}^* \mathring{\mathcal{N}}(0))$$

is moduli independent. □

Remark 2. The proof of the gluing formula holds verbatim for a more general class of metrics under the following two assumptions. First, the structure at infinity should be given by a finite union of conical/Euclidean ends. Second, we have to assume that nothing bad happens with the Laplace operator Δ^D of the compact part. In particular, it should have a well-defined zeta-function that extends to the complex plane with no pole at zero. This works for instance if the metric is smooth in the compact part or, if it is flat with conical singularities and the Friedrichs extension is chosen.

2.5. Closing the Euclidean (conical) ends with the help of gluing formulas. Let R be a sufficiently large positive number such that all the critical values of the meromorphic function f lie in the ball $\{|z| < R\}$.

In the holomorphic local parameter $\eta_j = y^{-1/k_j}$ in a vicinity U_j ($|y_j| > R$) of the j -th conical end of the angle $2\pi k_j$ ($k_j \geq 1$) of the Riemannian manifold $(X, |df|^2)$ (i.e., a pole of f of order k_j) the metric $\mathbf{m} = |df|^2$ takes the form

$$\mathbf{m} = k_j^2 \frac{|d\eta_j|^2}{|\eta_j|^{2k_j+2}}.$$

Let χ_j be a smooth function on \mathbb{C} such that $\chi_j(\eta) = \chi_j(|\eta|)$, $|\chi_j(\eta)| \leq 1$, $\chi_j(\eta) = 0$ if $|\eta| > (R + 1)^{-1/k_j}$, $\chi_j(\eta) = 1$ if $|\eta| < (R + 2)^{-1/k_j}$. Introduce the metric $\tilde{\mathbf{m}}$ on X such that

$$\tilde{\mathbf{m}} = \begin{cases} \mathbf{m} & \text{for } |z| < R, \\ [1 + (|\eta_j|^{2k_j+2} - 1)\chi_j(\eta_j)]\mathbf{m} & \text{in } U_j. \end{cases}$$

Since the (Friedrichs extension of) the Laplace operator $\Delta^{\tilde{\mathbf{m}}}$ has discrete spectrum and the corresponding operator ζ -function is regular at $s = 0$ (see, e.g., [21] and the references therein), one can define the determinant $\det^* \Delta^{\tilde{\mathbf{m}}}$ via usual Ray-Singer zeta-regularization. Moreover, for this determinant the usual BFK gluing formula ([4], Theorem B^*) holds (under the condition that the contour cutting the surface X does not pass through the conical singularities of the metric $\tilde{\mathbf{m}}$). Applying this standard BFK gluing formula, we get

$$(2.46) \quad \ln \det^* \Delta^{\tilde{\mathbf{m}}} = \ln C_0 + \ln \det_{\zeta} \Delta^D + \ln \det_{\zeta}^* \mathcal{N} + \ln \det \Delta_{ext}^{\tilde{\mathbf{m}}},$$

where $\Delta_{ext}^{\tilde{\mathbf{m}}}$ is the operator of the Dirichlet problem for $\Delta^{\tilde{\mathbf{m}}}$ in the union $\cup_j U_j$. Using conformal invariance we see that \mathcal{N} is the same as in Theorem 1 and C_0 is a moduli independent constant ($C_0 = \frac{\text{Area}(X, \tilde{\mathbf{m}})}{\text{length}(\Sigma)}$).

Now equation (2.46) and Theorem 1 imply the following proposition.

Proposition 5. *The relative zeta-regularized determinant $\det_{\zeta}(\Delta, \mathring{\Delta})$ and the zeta-regularized determinant $\det_{\zeta}^* \Delta^{\tilde{\mathbf{m}}}$ have the same variations with respect to moduli, i.e., one has*

$$(2.47) \quad \partial_{z_k} \ln \det_{\zeta}(\Delta, \mathring{\Delta}) = \partial_{z_k} \ln \det_{\zeta}^* \Delta^{\tilde{\mathbf{m}}}$$

for $k = 1, \dots, M$.

Thus, the relative determinant of the Laplacian on a non-compact surface (X, \mathbf{m}) with conical points and conical/Euclidean ends can be studied via consideration of the zeta-regularized determinant of Laplacian on a compact surface $(X, \tilde{\mathbf{m}})$ with conical points. The latter surface is flat everywhere except the conical singularities (whose positions vary when one changes the moduli z_1, \dots, z_M) and smooth ends of non-zero curvature which remain unchanged.

In the next two sections we study some spectral properties of compact surfaces with conical points. The final goal is to derive the variational formulas for $\ln \det_{\zeta}^* \Delta^{\tilde{\mathbf{m}}}$.

3. S-MATRIX

In this section we introduce the so-called S -matrix and relate its behavior at $\lambda = 0$ with the Schiffer projective connection. The definition of the S -matrix originates in the general theory of boundary triplets (see [13], sect. 13) and, more

specifically, the general theory of self-adjoint extensions of elliptic operators in singular settings (see [35]). Here we will closely follow [15]. However, we should point out that some normalization constants in the latter reference are erroneous and that the normalization we use here is slightly different.

It is convenient to introduce the S -matrix in the following general setting.

3.1. General setting and normalizations. Let $(X, \tilde{\mathbf{m}})$ be a compact singular 2d Riemannian manifold (possibly with boundary). Let P be an interior point of X such that in a neighborhood V of it X is isometric to a neighborhood of the tip of the Euclidean cone of angle $2\ell\pi$.

We set $X_0 := X \setminus \{P\}$ and $X_\varepsilon := X \setminus B(p, \varepsilon)$. We also denote by γ_r the circle of radius r centered at P .

We will occasionally use several different ways of parametrizing V .

- Polar coordinates $(r, \theta) \in (0, r_{\max}) \times \mathbb{R}/2\ell\pi\mathbb{Z}$,
- Local complex coordinate z . The one-form dz is well defined on $V \setminus \{P\}$ and extends to V as a holomorphic one-form α with a zero of order $\ell - 1$ at P . Note that it may not have a global holomorphic extension to X .
- Distinguished complex parameter y such that $\alpha = \ell y^{\ell-1} dy$ near P .

We now want to consider the Laplace operator that is associated with $\tilde{\mathbf{m}}$. We assume that the set of singularities of $\tilde{\mathbf{m}}$ consists of a finite number of conical points (in particular, it may consist of a single point P). Let Δ be the Friedrichs extension of the Laplace operator on X with domain consisting of smooth functions that vanish near the singularities.

Remark 3. Actually the choice of extension (that is, the prescription of a certain asymptotic at singular points to functions from the domain of the self-adjoint extension) should be made at each singularity (see [15]). Here we really care only about the choice of extension at P , where we are choosing the Friedrichs extension (all the functions from the domain of this extension are bounded near P). At other singularities one can choose any other extension, not necessarily the Friedrichs one.

By definition we set $H^2(X)$ to be the domain of Δ and by $H^1(X)$ to be its form domain. We denote by Δ_0 the restriction of Δ to functions in $H^2(X)$ that vanish near p and by Δ_0^* its formal adjoint. By choice, the self-adjoint extension Δ corresponds to the Friedrichs extension of Δ_0 . We will also denote by $H_0^2(X) := \text{dom } \overline{\Delta_0}$. Near P , we have

$$\Delta_0^* = -4\partial_z\partial_{\bar{z}} = -4\left(\ell^2|y|^{2(\ell-1)}\right)^{-1}\partial_y\partial_{\bar{y}}.$$

Introduce a cut-off function ρ such that ρ has support in $r \leq r_{\max}$ and equals 1 near $r = 0$. Define the functions F^0 , F_ν^a , and F_ν^h via

$$\begin{aligned} F^0(z) &= c_0 \ln(z\bar{z})\rho(z) = c_0 \ln(r^2)\rho(z), \\ F_\nu^a(z) &= c_\nu \bar{z}^{-\nu}\rho(z), \quad \nu = \frac{k}{\ell}, \quad 0 < k < \ell, \\ F_\nu^h(z) &= c_\nu z^{-\nu}\rho(z), \quad \nu = \frac{k}{\ell}, \quad 0 < k < \ell, \end{aligned}$$

where

$$(3.48) \quad c_0 = \frac{1}{2\sqrt{\ell\pi}}, \quad c_\nu = \frac{1}{2\sqrt{\nu\ell\pi}}.$$

Remark 4. The indices a and h correspond to “antiholomorphic” and “holomorphic” behavior of the corresponding functions F at 0 . In all the formulas below the index ν runs through the set $\{k/\ell\}_{k=1,\dots,\ell-1}$, where $2\pi\ell$ is the conical angle at P .

Separating variables near P , it can be shown that any function in $\text{dom}(\Delta_0^*)$ admits the following expression (cf. [21], [31]):

$$\begin{aligned}
 (3.49) \quad u &= c_0\Lambda^0(u) + c_0\Lambda^{0,-}(u)F^0(z) \\
 &+ \sum_{\nu} c_{\nu}\Lambda_{\nu}^{h,-}(u)z^{-\nu} + c_{\nu}\Lambda_{\nu}^{a,-}(u)\bar{z}^{-\nu} \\
 &+ \sum_{\nu} c_{\nu}\Lambda_{\nu}^h(u)z^{\nu} + c_{\nu}\Lambda_{\nu}^a(u)\bar{z}^{-\nu} + u_0,
 \end{aligned}$$

where the Λ are linear functionals on $\text{dom}(\Delta_0^*)$ that vanish on H_0^2 and u_0 is in H_0^2 . Moreover, one has

$$\overline{\Lambda_{\nu}^a(u)} = \Lambda_{\nu}^h(\bar{u}).$$

For $u, v \in \text{dom}(\Delta^*)$ we define

$$\mathcal{G}(u, v) := \langle \Delta^*u, v \rangle - \langle u, \Delta^*v \rangle.$$

The Green formula implies

$$(3.50) \quad \mathcal{G}(u, v) = \lim_{\epsilon \rightarrow 0} \frac{2}{i} \int_{\gamma_{\epsilon}} \partial_z u \bar{v} dz + u \partial_{\bar{z}} \bar{v} d\bar{z},$$

where the circle γ_{ϵ} is positively oriented. Since c_0 and c_{ν} satisfy $\frac{2c_0^2}{i} \int_{\gamma_{\epsilon}} \frac{dz}{z} = 1$ and $\frac{2\nu c_{\nu}^2}{i} \int_{\gamma_{\epsilon}} \frac{dz}{z} = 1$, the asymptotics (3.49) and (3.1) imply that

$$\begin{aligned}
 \mathcal{G}(u, v) &= \Lambda^{0,-}(u)\overline{\Lambda^0(v)} - \Lambda^0(u)\overline{\Lambda^{0,-}(v)} \\
 &+ \sum_{\nu} \Lambda_{\nu}^{h,-}(u)\overline{\Lambda_{\nu}^a(v)} - \Lambda_{\nu}^a(u)\overline{\Lambda_{\nu}^{h,-}(v)} \\
 &+ \sum_{\nu} \Lambda_{\nu}^h(u)\overline{\Lambda_{\nu}^{a,-}(v)} - \Lambda_{\nu}^{a,-}(u)\overline{\Lambda_{\nu}^h(v)}.
 \end{aligned}$$

The domain of the Friedrichs extension of Δ is characterized by requiring that all the coefficients with the superscript “-” vanish; it follows from (3.1) that the linear functionals $\Lambda_{\nu}^{a,h,0}$ are continuous over $H^2(X)$ and supported at P . It can be proved that any linear functional Λ that is continuous on $H^2(X)$ and supported at P can be written as a linear combination of the linear functionals $\Lambda_0, \Lambda_{\nu}^a$, and Λ_{ν}^h . Finally, we notice that one has the following representation for the space $H_0^2(X)$:

$$H_0^2(X) = \{u \in H^2(X) : \Lambda_{\nu}^{\sharp}(u) = 0 \ \forall \nu, \sharp = a, h ; \Lambda^0(u) = 0\}.$$

3.2. Definition of the S -matrix. We follow [15], paying special attention to conjugations and normalizing constants.

In the following the symbols \sharp and \flat are to be substituted by $0, h$, or a . When the superscript is 0 the subscript ν is 0 , when it is a or h , $\nu = \frac{j}{\ell}$ where j ranges from 1 to $\ell - 1$.

We define

$$\begin{aligned}
 (3.51) \quad f_{\nu}^{\sharp}(\cdot; \lambda) &= (\Delta_0^* - \lambda)F_{\nu}^{\sharp}, \quad g_{\nu}^{\sharp}(\cdot; \lambda) = -(\Delta - \lambda)^{-1} f_{\nu}^{\sharp}(\cdot; \lambda), \\
 G_{\nu}^{\sharp}(\cdot; \lambda) &= F_{\nu}^{\sharp} + g_{\nu}^{\sharp}(\cdot; \lambda), \quad S_{\mu\nu}^{\sharp\flat} = \Lambda_{\mu}^{\sharp} \left(g_{\nu}^{\flat}(\cdot; \lambda) \right).
 \end{aligned}$$

Observe that by definition the g -functions belong to H^2 , which makes the latter definition consistent when seeing Λ_μ^\sharp as a linear functional over H^2 . Since Λ_μ^\sharp also makes sense as a linear functional over $\ker(\Delta^* - \lambda)$, we may also write

$$S_{\mu\nu}^{\sharp b} = \Lambda_\mu^\sharp \left(G_\nu^b(\cdot; \lambda) \right).$$

Remark 5. The functions F, f, g depend on the initial choice of ρ but the linear functionals Λ and the functions G are cut-off independent.

The S -matrix is defined by blocks :

$$(3.52) \quad S := \begin{pmatrix} S^{00} & S_{0\nu}^{0h} & S_{0\nu}^{0a} \\ S_{\mu 0}^{h0} & S_{\mu\nu}^{hh} & S_{\mu\nu}^{ha} \\ S_{\mu 0}^{a0} & S_{\mu\nu}^{ah} & S_{\mu\nu}^{aa} \end{pmatrix}.$$

Remark 6. If there are several conical points on the surface, then there are several ways to define the S -matrix depending on how many points we want to take into account. The S -matrix defined in [15] takes into account all the conical points whereas the one defined here deals only with the conical point P (even if there are other conical points on the surface). Thus the S -matrix that is constructed here is only a part of the one from [15].

Applying Green’s formula, we get

$$(3.53) \quad \begin{aligned} \Lambda_\nu^\sharp(u) &= \mathcal{G}(u, \overline{F^\sharp}) = \int (\Delta - \lambda)u \cdot F_\nu^\sharp - u \cdot f_\nu^\sharp(\cdot; \lambda) dS \\ &= \int (\Delta - \lambda)u \cdot F_\nu^\sharp + u \cdot (\Delta - \lambda)g_\nu^\sharp(\cdot; \lambda) dS = \int (\Delta - \lambda)u \cdot G_\nu^\sharp dS \end{aligned}$$

for any test function $u \in H^2(X)$ (here dS is the area element on X). We have used here that $\overline{F^{0,a,h}} = F^{0,h,a} \in \text{dom}(\Delta_0^*)$, u and g are in $H^2(X)$, and that Δ is real (i.e., commutes with complex conjugation) and self-adjoint.

Applying (3.53) to the g -functions gives the following alternative expressions for the S -matrix entries (we omit the dependence on λ):

$$(3.54) \quad S_{\mu\nu}^{\sharp b} = \int G_\mu^\sharp (\Delta - \lambda) g_\nu^b dS = - \int G_\mu^\sharp f_\nu^b dS.$$

Remark 7. The S -matrix allows the description of the elements of $\ker(\Delta_0^* - \lambda)$ in the following way. For any element $u \in \text{dom}(\Delta_0^*)$, denote by L^\pm the column of its coefficients $\Lambda_\nu^{\sharp,\pm}$ that describe the singular behavior of u near P . We have

$$(\Delta_0^* - \lambda)u = 0 \Leftrightarrow L^+ = S(\lambda)L^-.$$

This gives a (pure) formal analogy with a typical scattering situation. In our setting, any solution to the equation $(\Delta_0^* - \lambda)u = 0$ plays the role of scattered field, L^\pm is the “incoming” and “outgoing” parts, and the S -matrix is the “scattering” matrix.

4. BASIC PROPERTIES OF THE S -MATRIX

4.0.1. *Analyticity and complex conjugation.* From the analyticity of the resolvent we see that the S -matrix depends analytically on λ . Besides, the expression

$$S_{\mu\nu}^{\sharp b}(\lambda) = - \int G_\mu^\sharp(\cdot; \lambda) f_\nu^b(\cdot; \lambda) dS$$

and the fact that

$$\overline{f_\nu^{a,h}(\cdot, \lambda)} = f_\nu^{h,a}(\cdot; \bar{\lambda}), \quad \overline{G^{a,h}(\cdot; \lambda)} = G_\nu^{h,a}(\cdot; \bar{\lambda}),$$

lead to the following identities:

$$(4.55) \quad \overline{S^{hh}(\lambda)} = S^{aa}(\bar{\lambda}), \quad \overline{S^{ah}(\lambda)} = S^{ha}(\bar{\lambda}).$$

4.0.2. *Behavior for λ going to $-\infty$.* For $r > 0$ consider the equation

$$-u''(r) - \frac{1}{r}u'(r) + \frac{\nu^2}{r^2}u(r) = \lambda u(r).$$

Since $\nu \neq 0$, any solution to this equation has the following asymptotic behavior near zero:

$$u(r) = a_-r^{-\nu} + a_+r^\nu + o(r^\nu)$$

and the vector space of solutions that belongs to $L^2(rdr)$ is one-dimensional. We set $k_\nu(r; \lambda)$ to be the unique solution to this equation which is in $L^2(rdr)$ and that is normalized in such a way that

$$\begin{aligned} F_\nu^h(r, \theta) - k_\nu(r; \lambda) \exp(-i\nu\theta) &= O(r^\nu), \\ F_\nu^a(r, \theta) - k_\nu(r; \lambda) \exp(i\nu\theta) &= O(r^\nu), \end{aligned}$$

(i.e., we adjust the coefficient of $r^{-\nu}$ in k_ν^\sharp so that it coincides with the coefficient of F_ν^\sharp). By inserting a cut-off ρ , we define

$$\begin{aligned} K_\nu^h(r, \theta; \lambda) &:= k_\nu(r; \lambda) \exp(-i\nu\theta)\rho(r), \\ K_\nu^a(r, \theta; \lambda) &:= k_\nu(r; \lambda) \exp(i\nu\theta)\rho(r), \end{aligned}$$

as functions on X . We compute $R_\nu^\sharp(\cdot; \lambda) = (\Delta_0^* - \lambda) K_\nu^\sharp$, where $\sharp = a, h$.

Lemma 3. *For $\sharp = a, h$ we have*

$$G_\nu^\sharp(\cdot; \lambda) = K_\nu^\sharp - [\Delta - \lambda]^{-1} R_\nu^\sharp.$$

Proof. By construction it is straightforward that both sides of the equation are in $\ker(\Delta_0^* - \lambda)$ and by choice of normalization, both share the same singular behavior. \square

We denote by $\kappa_\nu(\lambda)$ the coefficient of r^ν in the asymptotic expansion of $k_\nu(\cdot; \lambda)$.

Corollary 2. *As $\Re\lambda$ goes to $-\infty$ we have*

$$\begin{aligned} S^{hh}(\lambda) &= O(|\lambda|^{-\infty}), \quad S^{aa}(\lambda) = O(|\lambda|^{-\infty}), \\ S^{ah}(\lambda) &= \text{diag}(\kappa_\nu(\lambda)) + O(|\lambda|^{-\infty}), \\ S^{ha}(\lambda) &= \text{diag}(\kappa_\nu(\lambda)) + O(|\lambda|^{-\infty}). \end{aligned}$$

Proof. The asymptotic expansion of Bessel functions implies that

$$\|R_\nu^\flat\|_{L^2} = O(|\lambda|^{-\infty}).$$

Therefore $\Lambda_\nu^\sharp([\Delta - \lambda]^{-1} R_\nu^\flat) = O(|\lambda|^{-\infty})$. Thus all entries of the S -matrix are given by $\Lambda_\mu^\sharp(K_\nu^\flat)$ up to $O(|\lambda|^{-\infty})$. The first term is seen to be 0 except for the diagonal terms in S^{ah} or S^{ha} for which it is $\kappa_\nu(\lambda)$. \square

Remark 8. A different proof is given in [15] using the heat kernel.

4.0.3. *Differentiation with respect to λ .* We denote by a dot the differentiation with respect to λ . Differentiating the defining equation (3.4) for g_ν^b , we find that

$$\dot{g}_\nu^b = (\Delta - \lambda)^{-1} G_\nu^b.$$

Thus, we get

$$(4.56) \quad \dot{S}_{\mu\nu}^{\sharp b} = \Lambda_\mu^\sharp \left(\dot{g}_\nu^b \right) = \int G_\mu^\sharp G_\nu^b dS.$$

From this relation we deduce the following proposition.

Proposition 6. *For any λ , $S^{aa}(\lambda)$ and $S^{hh}(\lambda)$ are symmetric matrices and*

$$(4.57) \quad {}^t S^{ah}(\lambda) = S^{ha}(\lambda).$$

Proof. The expression for $\dot{S}_{\mu\nu}^{\sharp b}$ yields that \dot{S}^{aa} and \dot{S}^{hh} are symmetric matrices. Since both tend to symmetric matrices (actually 0) as λ goes to $-\infty$, the first part of the claim follows. In the same way we obtain

$$\dot{S}_{\mu\nu}^{ah} - \dot{S}_{\nu\mu}^{ha} = 0.$$

Since S^{ah} and ${}^t S^{ha}$ tend to the same diagonal matrix as λ goes to $-\infty$, the second part of the claim also follows. □

Combining the identities (4.55) and (4.57) we conclude that S^{ah} is hermitian for λ real; actually it is an analytic family of hermitian matrices, meaning that $S^{ah}(\lambda) = (S^{ah}(\bar{\lambda}))^*$.

4.0.4. *Behavior for λ going to 0.* The matrix $S(\lambda)$ is well defined a priori only for λ in the resolvent set of Δ . However, it is always possible to define the function $f_\nu^\sharp(\cdot; \lambda = 0)$. Whenever $\sharp \neq 0$ the latter function is in the range of Δ . We can thus find solutions $g_\nu^\sharp(\cdot; 0)$ to the equation

$$\Delta g_\nu^* = -f_\nu^*(\cdot, 0).$$

The latter solutions are defined only up to the addition of a constant. It follows that the definition of $S_{\mu\nu}^{\sharp b}(0)$ makes sense for $\sharp \neq 0$, $b \neq 0$ and the following proposition holds.

Proposition 7. *For $\sharp \neq 0$ and $b \neq 0$ the matrix-valued function $\lambda \mapsto S^{\sharp b}(\lambda)$ extends holomorphically to a neighborhood of 0. Moreover, $S^{\sharp b}(0)$ depends only on the conformal class of \mathfrak{m} .*

Proof. For λ close to 0 we have

$$g_\nu^\sharp = \frac{1}{\lambda} \int_X f_\nu(\cdot, \lambda) dS + g_\nu^{\sharp, \perp}(\cdot, \lambda),$$

where $\int_X g_\nu^{\sharp, \perp} = 0$. Since $\lambda \mapsto \int_X f_\nu^\sharp(\lambda; 0) = 0$ is holomorphic and vanish at 0, we obtain that $\lambda \mapsto G_\nu^*$ can be holomorphically continued to a neighborhood of 0. The first statement follows. The second statement follows by remarking that G_ν^\sharp is a function in $\text{dom}(\Delta_0^*)$ such that $\Delta_0^* G_\nu^\sharp = 0$ and the singular behavior near P is prescribed. Both conditions are conformally invariant so that if we change the metric in its conformal class, we may only change G by adding a constant. This will not affect the coefficients in the S -matrix we are considering here. □

4.1. **$S(0)$ and the Schiffer projective connection.** Choose a marking for the Riemann surface X , i.e., a symplectic set of generators $a_1, b_1, \dots, a_g, b_g$ of $H_1(X, \mathbf{Z})$. Let $\{v_1, \dots, v_g\}$ be the basis of holomorphic differentials on X normalized via

$$\int_{a_i} v_j = \delta_{ij}.$$

Then the matrix of b -periods of the marked Riemann surface X is defined via

$$\mathbf{B} = \left\| \int_{b_i} v_j \right\|.$$

Let $W(\cdot, \cdot)$ be the canonical meromorphic bidifferential on $X \times X$, with properties $W(P, Q) = W(Q, P)$,

$$\int_{a_i} W(\cdot, P) = 0,$$

and

$$\int_{b_j} W(\cdot, P) = 2\pi i v_j(P).$$

The bidifferential W has only a double pole along the diagonal $P = Q$. In any holomorphic local parameter $x(P)$ one has the asymptotics

$$(4.58) \quad W(x(P), x(Q)) = \left(\frac{1}{(x(P) - x(Q))^2} + H(x(P), x(Q)) \right) dx(P)dx(Q),$$

$$H(x(P), x(Q)) = \frac{1}{6} S_B(x(P)) + O(x(P) - x(Q)),$$

as $Q \rightarrow P$, where $S_B(\cdot)$ is the Bergman projective connection.

Consider the Schiffer bidifferential

$$\mathcal{S}(P, Q) = W(P, Q) - \pi \sum_{i,j} (\Im \mathbf{B})_{ij}^{-1} v_i(P) v_j(Q).$$

The Schiffer projective connection, S_{Sch} , is defined via the asymptotic expansion

$$\begin{aligned} \mathcal{S}(x(P), x(Q)) &= \left(\frac{1}{(x(P) - x(Q))^2} + \frac{1}{6} S_{Sch}(x(P)) + O(x(P) - x(Q)) \right) dx(P)dx(Q). \end{aligned}$$

One has the equality

$$(4.59) \quad S_{Sch}(x) = S_B(x) - 6\pi \sum_{i,j} (\Im \mathbf{B})_{ij}^{-1} v_i(x) \overline{v_j(x)}.$$

In contrast to the canonical meromorphic differential and the Bergman projective connection, the Schiffer bidifferential and the Schiffer projective connection are independent of the marking of the Riemann surface X .

Introduce also the so-called Bergman kernel (which is in fact the Bergman reproducing kernel for holomorphic differentials on X) as

$$B(x, \bar{x}) = \sum_{i,j} (\Im \mathbf{B})_{ij}^{-1} v_i(x) \overline{v_j(x)}.$$

Proposition 8. *Let X be a Riemann surface and let $\tilde{\mathbf{m}}$ be a conformal metric on X . Suppose that $\tilde{\mathbf{m}}$ has a conical singularity of angle $2\ell\pi$ at p . Let also x be the distinguished local parameter for $\tilde{\mathbf{m}}$ near p . Then there is the following relation between the entries of the holomorphic-holomorphic part, $S^{hh}(0)$, of the S -matrix:*

$$(4.60) \quad \sum_{k=1}^{\ell-1} \frac{\sqrt{k(\ell-k)}}{\ell} S_{\frac{k}{\ell} \frac{\ell-k}{\ell}}^{hh}(0) = -\frac{1}{6\ell(\ell-2)!} \left(\frac{d}{dx}\right)^{\ell-2} S_{Sch}(x) \Big|_{x=0}.$$

Remark 9. The same would hold true for a conical singularity of angle β with $2\pi(\ell-1) < \beta \leq 2\pi\ell$.

Remark 10. Observe that using the indices $\mu, \nu = \frac{k}{\ell}$ the left hand side of (4.60) can be written as

$$\sum_{\mu+\nu=1} \sqrt{\mu}\sqrt{\nu} S_{\mu\nu}^{hh}(0).$$

Proof. Introduce the following one-forms Ω_k and Σ_k on X :

$$\Omega_k = -\frac{1}{(k-1)!} \left(\frac{d}{dx}\right)^{k-1} \frac{W(\cdot, x)}{dx} \Big|_{x=0} + \frac{2\pi i}{(k-1)!} \sum_{\alpha, \beta} (\Im \mathbf{B})_{\alpha\beta}^{-1} \left\{ \Im v_{\beta}^{(k-1)}(0) \right\} v_{\alpha}(\cdot),$$

$$\Sigma_k = -i \frac{1}{(k-1)!} \left(\frac{d}{dx}\right)^{k-1} \frac{W(\cdot, x)}{dx} \Big|_{x=0} + \frac{2\pi i}{(k-1)!} \sum_{\alpha, \beta} (\Re \mathbf{B})_{\alpha\beta}^{-1} \left\{ \Re v_{\beta}^{(k-1)}(0) \right\} v_{\alpha}(\cdot),$$

where

$$v_{\beta}^{(k-1)}(0) := \left(\frac{d}{dx}\right)^{k-1} \frac{v_{\beta}(x)}{dx} \Big|_{x=0}.$$

All the periods of the differentials Ω_k and Σ_k are pure imaginary, therefore, one can correctly define the function f_k on X via

$$f_k(Q) = \Re \left\{ \int_{P_0}^Q \Omega_k \right\} - i \Re \left\{ \int_{P_0}^Q \Sigma_k \right\},$$

where P_0 is an arbitrary base point not coinciding with P . Clearly, f_k is harmonic in $X \setminus \{P\}$ and

$$(4.61) \quad f_k(x) = \frac{1}{x^k} + \text{const} + \sum_{j=1}^{\infty} (c_j x^j + d_j \bar{x}^j)$$

in a vicinity of P . One gets

$$c_l = -\frac{1}{l(k-1)!} \partial_x^{l-1} \partial_y^{k-1} H(x, y) \Big|_{x=y=0} + \frac{\pi}{l(k-1)!} \sum_{\alpha, \beta} (\Im \mathbf{B})_{\alpha\beta}^{-1} v_{\beta}^{(k-1)}(0) v_{\alpha}^{(l-1)}(0)$$

and

$$S_{\frac{k}{\ell} \frac{\ell-k}{\ell}}^{hh}(0) = \sqrt{\frac{\ell}{k}} c_{\ell}.$$

This implies that

$$\begin{aligned} \sum_{k=1}^{\ell-1} \frac{\sqrt{k(\ell-k)}}{\ell} S_{\frac{k}{\ell} \frac{\ell-k}{\ell}}^{hh}(0) &= -\frac{1}{\ell} \sum_{k=0}^{\ell-2} \frac{1}{k!(\ell-2-k)!} \partial_x^{\ell-2-k} \partial_y^k H(x, y) \Big|_{x=y=0} \\ &+ \frac{\pi}{\ell(\ell-2)!} \left(\frac{d}{dx}\right)^{\ell-2} \sum_{\alpha, \beta} (\Im \mathbf{B})_{\alpha\beta}^{-1} \frac{v_{\alpha}(x)v_{\beta}(x)}{(dx)^2} \Big|_{x=0}. \end{aligned}$$

Since

$$\frac{1}{6}S_B(x) = H(x, x) = \sum_{n=0}^{\infty} \frac{1}{n!} (\partial_x + \partial_y)^n H(x, y) \Big|_{x=y} x^n,$$

we have

$$\frac{1}{6}S_B^{(n)}(0) = \sum_{p=0}^n \frac{n!}{p!(n-p)!} \partial_x^p \partial_y^{n-p} H(0, 0),$$

which implies the proposition. □

Remark 11. From (4.61) with $k = 1$ it follows that for conical angles $2\pi < \beta \leq 4\pi$ we have

$$\begin{pmatrix} S^{hh}(0) & S^{ha}(0) \\ S^{ah}(0) & S^{aa}(0) \end{pmatrix} = \begin{pmatrix} -\frac{1}{6}S_{Sch}(0) & -\pi B(0, 0) \\ -\pi B(0, 0) & -\frac{1}{6}S_{Sch}(0) \end{pmatrix},$$

where the Schiffer projective connection and the Bergman kernel are calculated in the distinguished local parameter at P .

5. VARIATIONAL FORMULAS WITH RESPECT TO MODULI

In this section we derive the variational formulas for $\ln \det \Delta^{\tilde{\mathbf{m}}}$. This derivation goes as follows. First, using Kato-Rellich theory (see [18]), we prove variational formulas for the individual eigenvalues of the operator $\Delta^{\tilde{\mathbf{m}}}$. Using these formulas and the contour integral representation of the zeta-function of $\Delta^{\tilde{\mathbf{m}}}$, we express the variations of the value $\zeta'_{\Delta^{\tilde{\mathbf{m}}}}(0)$ with respect to the critical value z_k through a combination of the matrix elements of the S -matrix at the conical point P_k (the zero of the meromorphic differential df) of the metric $\tilde{\mathbf{m}}$. The latter combination is the one appearing in Proposition 8 and can be expressed through the Schiffer projective connection.

5.1. Variational formula for eigenvalues of $\Delta^{\tilde{\mathbf{m}}}$.

Remark 12. In this section we will use w for the moduli parameter and on the surface we will use the complex parameter z and (x, y) for the associated local cartesian coordinates (so that $z = x + iy$). We warn the reader that in the rest of this paper we use z_i as the moduli parameters and x as a local complex parameter on X .

5.1.1. *Moving conical points.* Let $\tilde{\mathbf{m}}$ be a metric as constructed in §2.5. Let P be one of its conical points. We wish to define a metric $\tilde{\mathbf{m}}_w$ corresponding to the shift of P by $w \in \mathbb{C}$. The following makes this construction precise.

Define \tilde{X}_w to be the ℓ -fold covering of \mathbb{C} with one ramification point at w so that \tilde{X}_w can be identified with the Euclidean cone of total angle $2\ell\pi$.

Fix a cut-off function ρ ($\rho \in C_0^\infty(\mathbb{R})$; $\rho = 1$ near 0) and define a map ϕ_w from \mathbb{C} to itself by

$$\phi_w(z) = z + \rho(|z|)w.$$

For w small enough, this defines a family of smooth self-diffeomorphisms of \mathbb{C} . The cone \tilde{X}_0 can be obtained by gluing together ℓ copies of the plane after cutting along a fixed half-line d that emanates from the origin. The cone \tilde{X}_w can then be obtained by gluing ℓ copies of \mathbb{C} after cutting it along $\phi_w(d)$.

The function ϕ_w thus defines a family of smooth diffeomorphisms from \tilde{X}_0 onto \tilde{X}_w . Let the metric g_w on \tilde{X}_0 be the pull-back of the Euclidean metric on \tilde{X}_w by ϕ_w .

We write $w = a + ib$ and use the local cartesian coordinates $x + iy = z$ near P . For the metric $g_w = A(x, y; w)dx^2 + 2B(x, y; w)dxdy + C(x, y; w)dy^2$ we obtain the following expressions:

$$(5.62) \quad \begin{pmatrix} A & B \\ B & C \end{pmatrix} = {}^t D\phi_w D\phi_w, \quad D\phi_w = \begin{pmatrix} 1 + \frac{ax}{r}\rho'_1(r) & \frac{ay}{r}\rho'_1(r) \\ \frac{bx}{r}\rho'_1(r) & 1 + \frac{by}{r}\rho'_1(r) \end{pmatrix}.$$

It follows by direct verification that the coefficients of g_w are polynomials in a, b . Observe that g_w coincides with g_0 outside a ball centered at P so that g_w can be smoothly extended by any Riemannian metric that coincides with the Euclidean one in an annulus centered at P . This allows us to define a metric $\tilde{\mathbf{m}}_w$ on our given setting $X \equiv \tilde{X}_0$ that corresponds to some \tilde{X}_w that is obtained by fixing the exterior of a small ball centered at $P \in X$ and, in an even smaller ball, by shifting the conical point by w .

We denote by J_w the jacobian determinant of the metric $\tilde{\mathbf{m}}_w$ on X , by q_w the Dirichlet energy quadratic form associated with $\tilde{\mathbf{m}}_w$, and by n_w the Riemannian $L^2(X)$ scalar product on $(X, \tilde{\mathbf{m}}_w)$; i.e., we set (for a real u that is supported near P)

$$(5.63) \quad \begin{aligned} q_w(u) &= \int_{\tilde{X}_0} [C(\partial_x u)^2 - 2B\partial_x u\partial_y u + A(\partial_y u)^2] J_w^{-\frac{1}{2}} dx dy, \\ n_w(u) &= \int u^2 J_w^{\frac{1}{2}} dx dy. \end{aligned}$$

Observe that $q_w(u)$ and $n_w(u)$ do not depend on w for u supported away from P .

In order to apply spectral perturbation theory, we will need the first order variations of $q_w(u)$ and $n_w(u)$. We prove the relevant lemma below.

Lemma 4. *For any $\lambda \in \mathbb{C}$ and any $u \in H^1(X)$ we have*

$$(5.64) \quad \begin{aligned} [-\partial_w q + \lambda \partial_w n]_{w=0}(u) &= 2 \int_{X_0} (\partial_z u)^2 \frac{z\rho'(r)}{r} dx dy + \frac{\lambda}{2} \int_{X_0} u^2 \frac{\bar{z}\rho'(r)}{r} dx dy, \\ [-\partial_{\bar{w}} q + \lambda \partial_{\bar{w}} n]_{w=0}(u) &= 2 \int_{X_0} (\partial_{\bar{z}} u)^2 \frac{\bar{z}\rho'(r)}{r} dx dy + \frac{\lambda}{2} \int_{X_0} u^2 \frac{z\rho'(r)}{r} dx dy. \end{aligned}$$

Proof. Denote by

$$\mathbb{G}_w := \begin{pmatrix} C & -B \\ -B & A \end{pmatrix}$$

so that we have

$$q_w(u) := \int_{X_0} {}^t \nabla u \mathbb{G} \nabla u \cdot J_w^{-\frac{1}{2}} dx dy.$$

Differentiating at $w = 0$, we obtain

$$\partial_w q_w(u) = \int_{X_0} {}^t \nabla u \cdot \left(\partial_w \mathbb{G} - \frac{1}{2} \partial_w J \mathbb{I} \right) \cdot \nabla u dS.$$

A straightforward computation yields

$$\begin{aligned} \partial_a \mathbb{G} - \frac{1}{2} \partial_a J \mathbb{I} &= \begin{pmatrix} -\frac{x\rho'}{r} & -\frac{y\rho'}{r} \\ -\frac{y\rho'}{r} & \frac{x\rho'}{r} \end{pmatrix}, \\ \partial_b \mathbb{G} - \frac{1}{2} \partial_b J \mathbb{I} &= \begin{pmatrix} \frac{y\rho'}{r} & -\frac{x\rho'}{r} \\ -\frac{x\rho'}{r} & -\frac{y\rho'}{r} \end{pmatrix}. \end{aligned}$$

From this we find

$$\begin{aligned} {}^t\nabla u \cdot \left(\partial_w \mathbb{G} - \frac{1}{2} \partial_w \mathbb{H} \right) \cdot \nabla u &= {}^t\nabla u \cdot \begin{pmatrix} -\frac{z\rho'}{2r} & \frac{iz\rho'}{2r} \\ \frac{iz\rho'}{2r} & \frac{z\rho'}{2r} \end{pmatrix} \cdot \nabla u \\ &= -\frac{z\rho'}{2r} ((\partial_x u)^2 - (\partial_y u)^2 - 2i\partial_x u \partial_y u) \\ &= -\frac{2z\rho'}{r} (\partial_z u)^2. \end{aligned}$$

The other terms proceed in the same way. □

5.1.2. *Variational formulas for eigenvalues of $\tilde{\mathbf{m}}$.* In this section, we compute variational formulas for the eigenvalues of $\tilde{\mathbf{m}}_w$. In order to do so we use the Kato-Rellich analytic perturbation theory [18]. It should be noticed that the family of metrics g_w is smooth in w but *not analytic*; see (5.62). We thus fix w and introduce $q_t = q_{tw}$ and $n_t = n_{tw}$, which are analytic in t . Because of this we may only consider directional derivatives.

The eigenvalue equation that gives the spectrum of q_t relatively to n_t is

$$(5.65) \quad q_t(u_t, v) = \lambda_t n_t(u_t, v).$$

This problem is analytic in t so that the eigenvalues are organized into real-analytic branches; see [18, Chapter VII.6.5] for details.

The first order variation for the eigenbranch (λ_t, u_t) is given by the following Feynman-Hellmann formula:

$$(5.66) \quad \frac{d\lambda}{dt} = \frac{dq}{dt}(u) - \lambda \frac{dn}{dt}(u)$$

which is obtained by differentiating equation (5.65) with v fixed and then evaluating at $v = u_t$.

Proposition 9. *Let r be small enough; then for $\lambda \in \text{spec}(\Delta^{\tilde{\mathbf{m}}})$ we have*

$$(5.67) \quad \begin{aligned} \partial_w \lambda &= \frac{2}{i} \int_{\gamma_r} (\partial_z u)^2 dz - \frac{\lambda}{4} u^2 d\bar{z}, \\ \partial_{\bar{w}} \lambda &= -\frac{2}{i} \int_{\gamma_r} (\partial_{\bar{z}} u)^2 d\bar{z} - \frac{\lambda}{4} u^2 dz. \end{aligned}$$

Let (λ_t, u_t) be an eigenbranch of q_t relatively to n_t ; then $\lambda' = \frac{d}{dt}|_{t=0} \lambda$ is given by

$$\lambda' = w \partial_w \lambda + \bar{w} \partial_{\bar{w}} \lambda,$$

where in the expression of $\partial_w \lambda$ and $\partial_{\bar{w}} \lambda$, $u = u_0$ is the eigenvector of the eigenbranch (λ_t, u_t) at $t = 0$.

Remark 13. We remind the reader of one subtlety of perturbation theory (see [18], [30]). In case of a multiple eigenvalue λ_0 , for any family q_t there are several eigenbranches emanating from λ_0 , and the corresponding initial eigenvectors may actually depend on the chosen family. In particular the expressions $\partial_w \lambda$ and $\partial_{\bar{w}} \lambda$ also depend on the initial w that defines $q_t := q_{tw}$. In other terms, for any direction w it is possible to organize the spectrum into eigenvalue branches but it may not be possible to organize the eigenvalues as functions that are differentiable with respect to w varying in the ball.

Proof. We start with the one-form

$$\omega_u = \rho(z) \cdot \left((\partial_z u)^2 dz - \frac{\lambda}{4} u^2 d\bar{z} \right).$$

Since (λ, u) is an eigenpair of the Laplace operator we compute

$$d\omega_u = - \left[\frac{z\rho'}{2r} (\partial_z u)^2 + \frac{\lambda}{4} \cdot \frac{\bar{z}\rho'}{2r} u^2 \right] dz \wedge d\bar{z}.$$

We now use the Stokes formula to obtain

$$\begin{aligned} \int_{\gamma_r} \omega_u &= - \int_X d\omega_u \\ &= \int_X \left[\frac{z\rho'}{2r} (\partial_z u)^2 + \frac{\lambda}{4} \cdot \frac{\bar{z}\rho'}{2r} u^2 \right] dz \wedge d\bar{z} \\ &= \frac{1}{2} \int_X \left[\frac{z\rho'}{r} (\partial_z u)^2 + \frac{\lambda}{4} \cdot \frac{\bar{z}\rho'}{r} u^2 \right] (-2idxdy). \end{aligned}$$

On the other hand, in (5.66) we use the formulas provided by Lemma 4 to obtain :

$$(5.68) \quad \partial_w \lambda = -2 \int_{X_0} (\partial_z u)^2 \frac{z\rho'(r)}{r} dx dy - \frac{\lambda}{2} \int_{X_0} u^2 \frac{\bar{z}\rho'(r)}{r} dx dy.$$

Comparing the two yields the first formula in (5.67). The second one follows either from the same computation or by complex conjugation. \square

Since ω_u is closed in $B(p, r_0) \setminus \{p\}$ we may tend to let r to 0 in the preceding formulas. We thus obtained a formula for $\partial_w \lambda$ that is expressed only through the asymptotic expansion of u near p .

Recall that by definition of the linear functionals Λ_ν^\sharp , we have in the local coordinate z the following expansion near p :

$$u(z) := c_0 \Lambda^0(u) + \sum_\nu c_\nu \Lambda_\nu^h(u) z^\nu + c_\nu \Lambda^a(u) \bar{z}^\nu + u_0$$

with $u_0 \in H_0^2$. By letting r go to zero we obtain the following lemma.

Lemma 5. *Let $A = [a_{\mu\nu}]$ be the matrix defined by*

$$\begin{cases} a_{\mu\nu} = 4\pi\mu c_\mu \cdot \nu c_\nu & \text{if } \mu + \nu = 1, \\ a_{\mu\nu} = 0 & \text{otherwise.} \end{cases}$$

We have the alternative expressions

$$(5.69) \quad \partial_w \lambda = \sum_{\mu+\nu=1} \Lambda_\nu^h(u) a_{\mu\nu} \Lambda_\mu^h(u), \quad \partial_{\bar{w}} \lambda = \sum_{\mu+\nu=1} \Lambda_\nu^a(u) a_{\mu\nu} \Lambda_\mu^a(u).$$

Proof. We prove the formula for $\partial_w \lambda$; the proof is the same for $\partial_{\bar{w}} \lambda$. First observe that since u is bounded we have

$$\lim_{r \rightarrow 0} \int_{\gamma_r} u^2 d\bar{z} = 0.$$

Now, if u_0 is smooth and compactly supported away from p , using the Stokes formula, we have that for any $v \in H^2$,

$$\int_{\gamma_r} \partial_z v \partial_z u_0 dz = \frac{1}{4} \int_{B_r} \Delta v \partial_z u_0 + \partial_z v \Delta u_0 dz \wedge d\bar{z}.$$

By continuity, this equality persists for $u_0 \in H_0^2$. It follows that for any $u \in H^2$ and any $u_0 \in H_0^2$ we have

$$\lim_{r \rightarrow 0} \int_{\gamma_r} \partial_z u \partial_{\bar{z}} u_0 dz = 0.$$

It follows that

$$\partial_w \lambda = \lim_{r \rightarrow 0} \int_{\gamma_r} (\partial_z(u - u_0))^2 dz.$$

By definition we have

$$u - u_0 = c_0 \Lambda^0(u) + \sum_{\nu} c_{\nu} \Lambda_{\nu}^h(u) z^{\nu} + c_{\nu} \Lambda^a(u) \bar{z}^{\nu},$$

so that the claim follows by a direct computation. □

Using the definition of Λ_{ν}^h and the fact that u is an eigenfunction, we obtain

Corollary 3. *For any $\lambda \in \mathbb{C} \setminus [0, \infty)$ the series $\sum_{\lambda_n \in \text{spec}(\Delta^{\tilde{m}})} \partial_w \lambda_n (\lambda_n - \lambda)^{-2}$ is absolutely convergent and*

$$\sum_{\lambda_n \in \text{spec}(\Delta^{\tilde{m}})} \frac{\partial_w \lambda_n}{(\lambda_n - \lambda)^2} = \text{Tr} \left(A \frac{\partial S^{hh}}{\partial \lambda}(\lambda) \right).$$

Proof. To prove the absolute convergence it suffices to show that for any ν we have

$$\sum_{\lambda_n \in \text{spec}(\Delta^{\tilde{m}})} \left| \frac{\Lambda_{\nu}^h(u_n)}{\lambda_n - \lambda} \right|^2 < \infty.$$

Since

$$\Lambda_{\nu}^h(u_n) = \int (\Delta - \lambda) u_n G_{\nu}^h = (\lambda_n - \lambda) \langle G_{\nu}^h, u_n \rangle,$$

the claim follows by remarking that the eigenfunctions u_n form an orthonormal basis. By the Plancherel formula we then obtain

$$\sum_{\lambda_n \in \text{spec}(\Delta^{\tilde{m}})} \frac{\partial_w \lambda_n}{(\lambda_n - \lambda)^2} = \sum_{\mu, \nu} a_{\mu\nu} \int_X G_{\mu}^h(x; \lambda) G_{\nu}^h(x; \lambda) dS.$$

We now remark that using (4.56)

$$\int_X G_{\mu}^h(x; \lambda) G_{\nu}^h(x; \lambda) dS = \frac{\partial_{\lambda} S_{\mu\nu}^{hh}}{\partial \lambda},$$

and A is a symmetric matrix. □

Remark 14. For $\sum_{\lambda_n \in \text{spec}(\Delta^{\tilde{m}})} \partial_{\bar{w}} \lambda_n (\lambda_n - \lambda)^{-2}$ we also have a similar formula involving S^{aa} .

5.2. Variational formula for $\zeta'(0; \Delta^{\tilde{m}})$. We prove the following proposition.

Proposition 10. *Let \tilde{m}_w be the family of metrics defined above, let S^{hh} be the holomorphic-holomorphic part of the corresponding S -matrix, and let A be the matrix defined in Lemma 5. We have*

$$(5.70) \quad \partial_w \zeta'(0; \Delta^{\tilde{m}}) = \text{Tr} (AS^{hh}(0)) = \sum_{\mu+\nu=1} \sqrt{\mu} \sqrt{\nu} S_{\mu\nu}^{hh}(0).$$

Proof. We start from the following integral representation of the zeta-function of the operator $\Delta^{\tilde{m}} - \lambda$ through the trace of the second power of the resolvent:

$$(5.71) \quad s\zeta(s + 1; \Delta^{\tilde{m}} - \lambda) = \frac{1}{2\pi i} \int_{\Gamma_\lambda} (z - \lambda)^{-s} \text{Tr}((\Delta^{\tilde{m}} - z)^{-2}) dz,$$

where Γ_λ is a contour connecting $-\infty + i\epsilon$ with $-\infty - i\epsilon$ and following the cut $(-\infty, \lambda)$ at the (sufficiently small) distance $\epsilon > 0$. Using Corollary 3, differentiation under the integral sign is legitimate and we obtain

$$(5.72) \quad s\partial_w \zeta(s + 1, \Delta^{\tilde{m}} - \lambda) = \frac{1}{2\pi i} \int_{\Gamma_\lambda} (z - \lambda)^{-s} \sum_{\lambda_n \in \text{spec}(\Delta^{\tilde{m}})} \frac{-2\partial_w \lambda_n}{(\lambda_n - z)^3} dz.$$

Using again Corollary 3, it is legitimate to integrate by parts to get

$$(5.73) \quad s\partial_w \zeta(s + 1, \Delta^{\tilde{m}} - \lambda) = \frac{-s}{2\pi i} \int_{\Gamma_\lambda} (z - \lambda)^{-s-1} \sum_{\lambda_n \in \text{spec}(\Delta^{\tilde{m}})} \frac{\partial_w \lambda_n}{(\lambda_n - z)^2} dz.$$

We can now divide by s , use Corollary 3 once again, and replace $s + 1$ by s to finally obtain

$$(5.74) \quad \partial_w \zeta(s, \Delta^{\tilde{m}} - \lambda) = \frac{-1}{2\pi i} \int_{\Gamma_\lambda} (z - \lambda)^{-s} \text{Tr}(A\partial_z S^{hh}(z)) dz.$$

Using the behavior of S^{hh} at infinity we can make an integration by parts again and obtain

$$(5.75) \quad \partial_w \zeta(s, \Delta^{\tilde{m}} - \lambda) = \frac{-s}{2\pi i} \int_{\Gamma_\lambda} (z - \lambda)^{-s-1} \text{Tr}(AS^{hh}(z)) dz.$$

Differentiating with respect to s and setting $s = 0$ gives

$$\partial_w \zeta'(0, \Delta^{\tilde{m}} - \lambda) = \frac{-1}{2\pi i} \int_{\Gamma_\lambda} (z - \lambda)^{-1} \text{Tr}(AS^{hh}(z)) dz.$$

The claim follows by applying Cauchy’s theorem. □

Now, using Proposition 5, the preceding proposition and Proposition 8 we arrive at the following corollary.

Corollary 4. *Let P_m be a zero of the meromorphic differential df of multiplicity ℓ_m and let $z_m = f(P_m)$ be the corresponding critical value of f . Let also $x_m = (z - z_m)^{\frac{1}{\ell_m+1}}$ be the distinguished local parameter in a vicinity of P_m . Then*

$$(5.76) \quad \partial_{z_m} \ln \det_\zeta(\Delta, \mathring{\Delta}) = \frac{1}{6(\ell_m + 1)(\ell_m - 1)!} \left(\frac{d}{dx_m} \right)^{\ell_m-1} S_{Sch}(x_m) \Big|_{x_m=0}.$$

6. INTEGRATION OF THE EQUATIONS FOR $\ln \det$ AND EXPLICIT EXPRESSIONS FOR THE τ -FUNCTION

Let, as before, \mathbb{B} be the matrix of b -periods of the Torelli marked Riemann surface X and let $\{v_\alpha\}_{\alpha=1,\dots,g}$ be the basis of the normalized holomorphic differentials on X . Using the Rauch formulas (see, e.g., [24], [25], [22]),

$$\partial_{z_m} \mathbb{B}_{\alpha\beta} = \oint_{P_m} \frac{v_\alpha v_\beta}{df},$$

one immediately gets the relation

$$(6.77) \quad \partial_{z_m} \ln \det \mathfrak{S}\mathbb{B} = \frac{1}{2i} \text{Tr} [(\partial_{z_m} \mathbb{B})(\mathfrak{S}\mathbb{B})^{-1}] = \frac{1}{2i} \oint_{P_m} \frac{\sum_{\alpha\beta} \mathfrak{S}\mathbb{B}_{\alpha\beta}^{-1} v_\alpha v_\beta}{df},$$

where the contour integrals are taken over a small contour on X encircling the point P_m (in the positive direction).

Now, using the relation (4.59), the equations (6.77) and (5.76) together with elementary properties of the Schwarzian derivative (see, e.g., [39]), we arrive at the following version of Corollary 4 rewritten in the invariant form.

Theorem 5. *Let P_m be a zero of the meromorphic differential df of multiplicity ℓ_m and let $z_m = f(P_m)$ be the corresponding critical value of f . Let also $x_m = (z - z_m)^{\frac{1}{\ell_m+1}}$ be the distinguished local parameter in a vicinity of P_m . Then*

$$(6.78) \quad \partial_{z_m} \ln \frac{\det_\zeta(\Delta, \hat{\Delta})}{\det \mathfrak{S}\mathbb{B}} = -\frac{1}{12\pi i} \oint_{P_m} \frac{S_B - S_f}{df},$$

where S_B is the Bergman projective connection and $S_f = \frac{f'''f' - \frac{3}{2}(f'')^2}{(f')^2}$ is the Schwarzian derivative.

Remark 15. Notice that the difference $S_B - S_f$ is a quadratic differential and, therefore, the integrand in (6.78) is a meromorphic one-form.

Equation (6.78) implies the relation

$$(6.79) \quad \det_\zeta(\Delta, \hat{\Delta}) = C \det \mathfrak{S}\mathbb{B} |\tau|^2,$$

where τ is a holomorphic function of moduli z_1, \dots, z_M (actually, τ is a holomorphic section of some holomorphic line bundle over the Hurwitz space, see [27] for further information; here we restrict ourselves to local considerations: the reader may assume for simplicity that everything happens in a small vicinity of the covering $f : X \rightarrow \mathbb{C}P^1$ in the Hurwitz space $\mathcal{H}(M, N)$) subject to the system of PDE

$$(6.80) \quad \partial_{z_m} \ln \tau = -\frac{1}{12\pi i} \oint_{P_m} \frac{S_B - S_f}{df}$$

and C is a moduli independent constant.

The system of PDE (6.80) first appeared in the context of the theory of isomonodromic deformations and Frobenius manifolds in [22] and [24], where, in particular, it was explicitly integrated. We recall these results in the next subsection.

6.1. Explicit expressions for τ . In this section we recall explicit formulas for the holomorphic solution, τ , of the system (6.80) derived in [24], [22] (see also [26] and [25] for alternative and more straightforward proofs). The result should be formulated separately for low genera $g = 0, 1$ and for higher genus $g > 1$. We start with the higher genus situation.

Let $g > 1$. Take a non-singular odd theta characteristic δ and consider the corresponding theta-function $\theta[\delta](t; \mathbb{B})$, where $t = (t_1, \dots, t_g) \in \mathbb{C}^g$. Put

$$\omega_\delta = \sum_{i=1}^g \frac{\partial \theta[\delta]}{\partial t_i} (0; \mathbb{B}) \omega_i.$$

All zeros of the holomorphic 1-differential ω_δ have even multiplicities, and $\sqrt{\omega_\delta}$ is a well-defined holomorphic spinor on X . Following Fay [9], consider the prime form

$$(6.81) \quad E(x, y) = \frac{\theta[\delta] \left(\int_x^y v_1, \dots, \int_x^y v_g; \mathbb{B} \right)}{\sqrt{\omega_\delta(x)} \sqrt{\omega_\delta(y)}}.$$

To make the integrals uniquely defined, we fix $2g$ simple closed loops in the homology classes a_i, b_i that cut X into a connected domain, and pick the integration paths that do not intersect the cuts. The sign of the square root is chosen so that

$$E(x, y) = \frac{\zeta(y) - \zeta(x)}{\sqrt{d\zeta(x)} \sqrt{d\zeta(y)}} (1 + O((\zeta(y) - \zeta(x))^2))$$

as $y \rightarrow x$, where ζ is a local parameter such that $d\zeta = \omega_\delta$.

We introduce local coordinates on X that we call *distinguished* with respect to f . Consider the divisor $(df) = \sum_k d_k p_k$, $p_k \in X$, $d_k \in \mathbb{Z}, d_k \neq 0$, of the meromorphic differential df . We take $z = f(x)$ as a local coordinate on $X - \bigcup_k p_k$ and

$$(6.82) \quad x_k = \begin{cases} (f(x) - f(p_k))^{\frac{1}{d_k+1}} & \text{if } d_k > 0, \\ f(x)^{\frac{1}{d_k+1}} & \text{if } d_k < 0, \end{cases}$$

near $p_k \in X$. In terms of these coordinates we have $E(x, y) = \frac{E(z(x), z(y))}{\sqrt{dz(x)} \sqrt{dz(y)}}$, and we define

$$\begin{aligned} E(z, p_k) &= \lim_{y \rightarrow p_k} E(z(x), z(y)) \sqrt{\frac{dz_k}{dz}}(y), \\ E(p_k, p_l) &= \lim_{\substack{x \rightarrow p_k \\ y \rightarrow p_l}} E(z(x), z(y)) \sqrt{\frac{dx_k}{d\zeta}}(x) \sqrt{\frac{dx_l}{d\zeta}}(y). \end{aligned}$$

Let \mathcal{A}^x be the Abel map with the basepoint x , and let $K^x = (K_1^x, \dots, K_g^x)$ be the vector of Riemann constants

$$(6.83) \quad K_i^x = \frac{1}{2} + \frac{1}{2} \mathbb{B}_{ii} - \sum_{j \neq i} \int_{a_i} \int_x^y v_j \, dy$$

(as above, we assume that the integration paths do not intersect the cuts on X). Then we have $\mathcal{A}^x((df)) + 2K^x = \Omega Z + Z'$ for some $Z, Z' \in \mathbb{Z}^g$. One has the following expression for the holomorphic solution to (6.80)(see [24], here we follow the presentation of this result in [27]):

$$(6.84) \quad \tau(X, f) = \frac{\left(\left(\sum_{i=1}^g v_i(\zeta) \frac{\partial}{\partial t_i} \right)^g \theta(t; \mathbb{B}) \Big|_{t=K\zeta} \right)^{2/3}}{e^{6^{-1}\pi\sqrt{-1}(\mathbb{B}Z+4K\zeta, Z)} W(\zeta)^{2/3}} \frac{\prod_{k < l} E(p_k, p_l)^{6^{-1}d_k d_l}}{\prod_k E(\zeta, p_k)^{3^{-1}(g-1)d_k}}.$$

Here $\theta(t; \mathbb{B}) = \theta[0](t; \mathbb{B})$ is the Riemann theta-function, $t = (t_1, \dots, t_g) \in \mathbb{C}^g$, and W is the Wronskian of the normalized holomorphic differentials v_1, \dots, v_g on X ; the expression in (6.84) is independent of $\zeta \in X$.

Let $g = 1$. Then the function $\tau(X, f)$ is given (see [22]) by the equation

$$(6.85) \quad \tau(X, f) = [\theta_1'(0 | \mathbb{B})]^{2/3} \frac{\prod_{k=1}^K h_j^{(k_j+1)/12}}{\prod_{m=1}^M f_m^{\ell_m/12}},$$

where $v(P)$ is the normalized Abelian differential on the elliptic Torelli marked curve X ; $v(P) = f_m(x_m)dx_m$ near P_m , where $x_m = (z - z_m)^{1/(\ell_m+1)}$ is the distinguished

local parameter near the zero, P_m of the differential df ; $f_m \equiv f_m(0)$; $v(P) = h_j(\zeta_j)d\zeta_j$ as $P \rightarrow \infty_j$, $\zeta_j = z^{-1/k_j}$, where k_j is the multiplicity of the pole ∞_j of f , $h_k \equiv h_k(0)$; θ_1 is the jacobi theta-function.

Let $g = 0$ and let $U : X \rightarrow \mathbb{P}^1$ be a biholomorphic map such that $U(\infty_1) = \infty$ and $U(P) = (f(P))^{1/k_1} + o(1)$ as $P \rightarrow \infty_1$. Then (see [26])

$$(6.86) \quad \tau(X, f) = \frac{\prod_{j=2}^K (\frac{dU}{d\zeta_j} |_{\zeta_j=0})^{(k_j+1)/12}}{\prod_{m=1}^M (\frac{dU}{dx_m} |_{x_m=0})^{l_m/12}}.$$

Summarizing (6.79) and (6.86), (6.85) (6.84), we get the main result of the present paper.

Theorem 6. *Let (X, f) be an element of the Hurwitz space $\mathcal{H}(M, N)$ and let $\tau(X, f)$ be given by expressions (6.86), (6.85), (6.84). There is the following explicit expression for the regularized relative determinant of the Laplacian Δ on the Riemann surface X :*

$$(6.87) \quad \det_{\zeta}(\Delta, \mathring{\Delta}) = C \det \mathfrak{S}\mathbb{B} |\tau|^2,$$

where C is a constant dependent only on the connected component of the space $\mathcal{H}(M, N)$ containing the element (X, f) .

6.2. Examples in genus 0. We finish the paper with two simple and especially instructive examples of the calculation of the determinant of the Laplacian Δ in genus 0.

Example 1. Let p be a polynomial with $N - 1$ simple critical points w_1, \dots, w_{N-1} and let the corresponding critical values be z_1, \dots, z_{N-1} (or, what is the same, a ramified covering with $N - 1$ simple branch points and one branch point of multiplicity N over the point at infinity of the base. In other words p is an element of the Hurwitz space $\mathcal{H}_{0,N}([1]^N)$ of meromorphic functions of degree N on the Riemann sphere \mathbb{P}^1 with a single pole of multiplicity N .

Let also w be the holomorphic coordinate on the cover \mathbb{P}^1 (more precisely, on $\mathbb{P}^1 \setminus \{\infty\}$) and let z be the holomorphic coordinate on the base \mathbb{P}^1 . One can assume that the leading coefficient of the polynomial $p(w)$ is equal to one.

Introduce the distinguished local parameter $x_k = \sqrt{z - z_k}$ at P_k . Then for $x_k \neq 0$ one has

$$\frac{dw}{dx_k} = \frac{1}{z'(w)} 2x_k = \frac{w - w_k}{p'(w) - p'(w_k)} \frac{2x_k}{w - w_k}.$$

Passing to the limit $x_k \rightarrow 0$, one gets

$$2 \left[w'(x_k) \Big|_{x_k=0} \right]^2 = \frac{1}{p''(w_k)}.$$

Thus,

$$(6.88) \quad \tau = \prod_{k=1}^{N-1} \left[w'(x_k) \Big|_{x_k=0} \right]^{-\frac{1}{12}} = \left\{ \prod_{k=1}^{N-1} p''(w_k) \right\}^{\frac{1}{24}} = \mathcal{R}(p', p'')^{\frac{1}{24}},$$

where $\mathcal{R}(f, g)$ is the resultant of polynomials f and g (since the τ -function is defined up to multiplicative constant, the power of 2 is omitted) and

$$\det_{\zeta}(\Delta^{|dp|^2}, \mathring{\Delta}) = C |\mathcal{R}(p', p'')|^{\frac{1}{12}}.$$

Example 2. Let r is a rational function with three simple poles (which can be assumed to coincide with $\infty, 0, 1$),

$$r(w) = aw - \frac{b}{w} - \frac{c}{w-1} + d;$$

i.e., r is an element of the Hurwitz space $\mathcal{H}_{0,3}(1, 1, 1)$ of meromorphic functions on P^1 of degree three with three simple poles.

Introducing the local parameter $\zeta = \frac{1}{z}$ in vicinities of the poles $w = 0$ and $w = 1$ of the cover, one gets for $\zeta \neq 0$

$$w'(\zeta) = -\frac{1}{r'(w)}r^2(w)$$

and, say, for $w = 1$

$$w'(0) = -\lim_{w \rightarrow 1} \frac{1}{r'(w)}r^2(w) = -c.$$

Analogously, $w'(0) = -b$ at the pole $w = 0$. For the local parameter $\tilde{w} = w/a$ (for which $\tilde{w}(P) = z(P) + o(1)$ as P tends to the pole $w = \infty$ of the cover) one has $\tilde{w}'(0) = -b/a$ at the pole $w = 0$ and $\tilde{w}'(0) = -c/a$ at the pole $w = 1$. On the other hand, writing $r'(w) = \frac{f}{g}$, where f and g are two polynomials and introducing the local parameter $x_k = \sqrt{z - z_k}$ near the critical point w_k of r ($k = 1, 2, 3, 4$), one gets similarly to (6.88)

$$\begin{aligned} \tilde{w}'(x_k) &= a^{-1}w'(x_k), \\ C \prod_{k=1}^4 [w'(x_k)|_{x_k=0}]^2 &= \prod_{k=1}^4 \frac{1}{r''(w_k)} = \frac{\mathfrak{R}(f, g)}{\mathfrak{R}(f, f')}. \end{aligned}$$

Calculating the resultants, one gets

$$\tau^{24} = a^3 b^3 c^3 \mathfrak{M}(a, b, c),$$

where

$$\mathfrak{M}(a, b, c) = a^3 + b^3 + c^3 + 3a^2b + 3a^2c + 3b^2a + 3b^2c + 3c^2a + 3c^2b - 21abc$$

and

$$\det_{\zeta}(\Delta^{|\mathrm{dr}|^2}, \mathring{\Delta}) = C|abc|^{1/4}|\mathfrak{M}(a, b, c)|^{1/12}.$$

ACKNOWLEDGMENTS

This work was finished during the stay of the third author at the Max Planck Institute of Mathematics (Bonn). The third author thanks the Institute for their hospitality and excellent working conditions. The work of the first author was partly supported by the ANR program Gerasic-ANR-13-BS01-0007-0. The work of the third author was supported by NSERC and ANR program Gerasic-ANR-13-BS01-0007-0.

REFERENCES

[1] Milton Abramowitz and Irene A. Stegun, *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, National Bureau of Standards Applied Mathematics Series, vol. 55, For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C., 1964. MR0167642

[2] Luis Alvarez-Gaumé, Jean-Benoît Bost, Gregory Moore, Philip Nelson, and Cumrun Vafa, *Bosonization on higher genus Riemann surfaces*, Comm. Math. Phys. **112** (1987), no. 3, 503–552. MR908551

- [3] M. Bershadsky and A. Radul, *g-loop amplitudes in bosonic string theory in terms of branch points*, Phys. Lett. B **193** (1987), no. 2-3, 213–218. MR900095
- [4] D. Burghelea, L. Friedlander, and T. Kappeler, *Meyer-Vietoris type formula for determinants of elliptic differential operators*, J. Funct. Anal. **107** (1992), no. 1, 34–65. MR1165865
- [5] Gilles Carron, *Déterminant relatif et la fonction ξ* (French, with French summary), Amer. J. Math. **124** (2002), no. 2, 307–352. MR1890995
- [6] Gilles Carron, *Le saut en zéro de la fonction de décalage spectral* (French, with English and French summaries), J. Funct. Anal. **212** (2004), no. 1, 222–260. MR2067165
- [7] Eric D’Hoker and D. H. Phong, *On determinants of Laplacians on Riemann surfaces*, Comm. Math. Phys. **104** (1986), no. 4, 537–545. MR841668
- [8] Boris Dubrovin, *Geometry of 2D topological field theories*, Integrable systems and quantum groups (Montecatini Terme, 1993), Lecture Notes in Math., vol. 1620, Springer, Berlin, 1996, pp. 120–348. MR1397274
- [9] John Fay, *Kernel functions, analytic torsion, and moduli spaces*, Mem. Amer. Math. Soc. **96** (1992), no. 464, vi+123. MR1146600
- [10] Robin Forman, *Functional determinants and geometry*, Invent. Math. **88** (1987), no. 3, 447–493. MR884797
- [11] William Fulton, *Hurwitz schemes and irreducibility of moduli of algebraic curves*, Ann. of Math. (2) **90** (1969), 542–575. MR0260752
- [12] Phillip Griffiths and Joseph Harris, *Principles of algebraic geometry*, Wiley-Interscience [John Wiley & Sons], New York, 1978. Pure and Applied Mathematics. MR507725
- [13] Gerd Grubb, *Distributions and operators*, Graduate Texts in Mathematics, vol. 252, Springer, New York, 2009. MR2453959
- [14] Andrew Hassell and Steve Zelditch, *Determinants of Laplacians in exterior domains*, Internat. Math. Res. Notices **18** (1999), 971–1004. MR1722360
- [15] Luc Hillairet and Alexey Kokotov, *Krein formula and S-matrix for Euclidean surfaces with conical singularities*, J. Geom. Anal. **23** (2013), no. 3, 1498–1529. MR3078362
- [16] Luc Hillairet, Victor Kalvin, and Alexey Kokotov, *Spectral determinants on Mandelstam diagrams*, Comm. Math. Phys. **343** (2016), no. 2, 563–600. MR3477347
- [17] M. V. Fedoryuk, *Metod perevala* (Russian), Izdat. “Nauka”, Moscow, 1977. MR0507923
- [18] Tosio Kato, *Perturbation theory for linear operators*, Classics in Mathematics, Springer-Verlag, Berlin, 1995. Reprint of the 1980 edition. MR1335452
- [19] Bernard S. Kay and Urban M. Studer, *Boundary conditions for quantum mechanics on cones and fields around cosmic strings*, Comm. Math. Phys. **139** (1991), no. 1, 103–139. MR1116412
- [20] V. G. Knizhnik, *Analytic fields on Riemann surfaces. II*, Comm. Math. Phys. **112** (1987), no. 4, 567–590. MR910579
- [21] Alexey Kokotov, *On the spectral theory of the Laplacian on compact polyhedral surfaces of arbitrary genus*, Computational approach to Riemann surfaces, Lecture Notes in Math., vol. 2013, Springer, Heidelberg, 2011, pp. 227–253. MR2920506
- [22] A. Kokotov and D. Korotkin, *Tau-functions on Hurwitz spaces*, Math. Phys. Anal. Geom. **7** (2004), no. 1, 47–96. MR2053591
- [23] Alexey Kokotov, *Polyhedral surfaces and determinant of Laplacian*, Proc. Amer. Math. Soc. **141** (2013), no. 2, 725–735. MR2996977
- [24] A. Kokotov and D. Korotkin, *Isomonodromic tau-function of Hurwitz Frobenius manifolds and its applications*, Int. Math. Res. Not. (2006), Art. ID 18746, 34. MR2211143
- [25] Aleksey Kokotov and Dmitry Korotkin, *Tau-functions on spaces of abelian differentials and higher genus generalizations of Ray-Singer formula*, J. Differential Geom. **82** (2009), no. 1, 35–100. MR2504770
- [26] Alexey Kokotov and Ian A. B. Strachan, *On the isomonodromic tau-function for the Hurwitz spaces of branched coverings of genus zero and one*, Math. Res. Lett. **12** (2005), no. 5-6, 857–875. MR2189245
- [27] A. Kokotov, D. Korotkin, and P. Zograf, *Isomonodromic tau function on the space of admissible covers*, Adv. Math. **227** (2011), no. 1, 586–600. MR2782203
- [28] Maxim Kontsevich and Simeon Vishik, *Geometry of determinants of elliptic operators*, Functional analysis on the eve of the 21st century, Vol. 1 (New Brunswick, NJ, 1993), Progr. Math., vol. 131, Birkhäuser Boston, Boston, MA, 1995, pp. 173–197. MR1373003

- [29] Maxim Kontsevich and Anton Zorich, *Connected components of the moduli spaces of Abelian differentials with prescribed singularities*, *Invent. Math.* **153** (2003), no. 3, 631–678. MR2000471
- [30] Andreas Kriegl and Peter W. Michor, *Differentiable perturbation of unbounded operators*, *Math. Ann.* **327** (2003), no. 1, 191–201. MR2006008
- [31] Edith A. Mooers, *Heat kernel asymptotics on manifolds with conic singularities*, *J. Anal. Math.* **78** (1999), 1–36. MR1714065
- [32] Werner Müller, *Relative zeta functions, relative determinants and scattering theory*, *Comm. Math. Phys.* **192** (1998), no. 2, 309–347. MR1617554
- [33] Werner Müller, *On the L^2 -index of Dirac operators on manifolds with corners of codimension two. I*, *J. Differential Geom.* **44** (1996), no. 1, 97–177. MR1420351
- [34] S. M. Natanzon, *Topology of 2-dimensional coverings and meromorphic functions on real and complex algebraic curves*, *Selecta Math. Soviet.* **12** (1993), no. 3, 251–291. Selected translations. MR1244839
- [35] Sergey A. Nazarov and Boris A. Plamenevsky, *Elliptic problems in domains with piecewise smooth boundaries*, *De Gruyter Expositions in Mathematics*, vol. 13, Walter de Gruyter & Co., Berlin, 1994. MR1283387
- [36] Peter Sarnak, *Extremal geometries*, *Extremal Riemann surfaces* (San Francisco, CA, 1995), *Contemp. Math.*, vol. 201, Amer. Math. Soc., Providence, RI, 1997, pp. 1–7. MR1429189
- [37] Barry Simon, *Notes on infinite determinants of Hilbert space operators*, *Advances in Math.* **24** (1977), no. 3, 244–273. MR0482328
- [38] Hidenori Sonoda, *Functional determinants on punctured Riemann surfaces and their application to string theory*, *Nuclear Phys. B* **294** (1987), no. 1, 157–192. MR909430
- [39] A. N. Tjurin, *Periods of quadratic differentials* (Russian), *Uspekhi Mat. Nauk* **33** (1978), no. 6(204), 149–195, 272. MR526014
- [40] D. R. Yafaev, *Matematicheskaya teoriya rasseyaniya* (Russian, with Russian summary), *Izdatel'stvo Sankt-Peterburgskogo Universiteta*, St. Petersburg, 1994. *Obshchaya teoriya*. [General theory]. MR1784870
- [41] Richard A. Wentworth, *Precise constants in bosonization formulas on Riemann surfaces. I*, *Comm. Math. Phys.* **282** (2008), no. 2, 339–355. MR2421480
- [42] Scott A. Wolpert, *Asymptotics of the spectrum and the Selberg zeta function on the space of Riemann surfaces*, *Comm. Math. Phys.* **112** (1987), no. 2, 283–315. MR905169
- [43] Al. B. Zamolodchikov, *Conformal scalar field on the hyperelliptic curve and critical Ashkin-Teller multipoint correlation functions*, *Nuclear Phys. B* **285** (1987), no. 3, 481–503. MR897030

MAPMO (UMR 7349 UNIVERSITÉ D'ORLÉANS-CNRS) UFR SCIENCES, BÂTIMENT DE MATHÉMATIQUES RUE DE CHARTRES, BP 6759 45067 ORLÉANS CEDEX 02, FRANCE

Email address: luc.hillairet@univ-orleans.fr

DEPARTMENT OF MATHEMATICS AND STATISTICS, CONCORDIA UNIVERSITY, 1455 DE MAISON-NEUVE BOULAVARD WEST, MONTREAL, QUEBEC, H3G 1M8 CANADA

Email address: vkalvin@gmail.com

DEPARTMENT OF MATHEMATICS AND STATISTICS, CONCORDIA UNIVERSITY, 1455 DE MAISON-NEUVE BOULAVARD WEST, MONTREAL, QUEBEC, H3G 1M8 CANADA

Email address: alexey.kokotov@concordia.ca